


Complete Genome Sequence of the Barley Pathogen *Xanthomonas translucens* pv. *translucens* DSM 18974^T (ATCC 19319^T)

Sebastian Jaenicke,^a Boyke Bunk,^b Daniel Wibberg,^c Cathrin Spröer,^b Lena Hersemann,^d Jochen Blom,^a Anika Winkler,^c Sarah Schatschneider,^{c*} Stefan P. Albaum,^c  Roland Kölliker,^d Alexander Goesmann,^a Alfred Pühler,^c Jörg Overmann,^b Frank-Jörg Vorhölter^{c*}

Justus-Liebig-University Giessen, Bioinformatics and Systems Biology, Giessen, Germany^a; Leibniz Institute DSMZ, German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany^b; Center for Biotechnology CeBiTec, Bielefeld University, Bielefeld, Germany^c; Agroscope, Molecular Ecology, Zurich, Switzerland^d

* Present address: Sarah Schatschneider, Evonik Nutrition and Care GmbH, Halle, Germany; Frank-Jörg Vorhölter, MVZ Dr. Eberhard and Partner Dortmund, Dortmund, Germany.

S.J. and B.B. contributed equally to this work.

We report here the complete 4.7-Mb genome sequence of *Xanthomonas translucens* pv. *translucens* DSM 18974^T, which causes black chaff disease on barley (*Hordeum vulgare*). Genome data of this *X. translucens* type strain will improve our understanding of this bacterial species.

Received 4 October 2016 Accepted 7 October 2016 Published 1 December 2016

Citation Jaenicke S, Bunk B, Wibberg D, Spröer C, Hersemann L, Blom J, Winkler A, Schatschneider S, Albaum SP, Kölliker R, Goesmann A, Pühler A, Overmann J, Vorhölter F-J. 2016. Complete genome sequence of the barley pathogen *Xanthomonas translucens* pv. *translucens* DSM 18974^T (ATCC 19319^T). *Genome Announc* 4(6):e01334-16. doi:10.1128/genomeA.01334-16.

Copyright © 2016 Jaenicke et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Frank-Jörg Vorhölter, frank@cebitec.uni-bielefeld.de.

The Gram-negative gammaproteobacterium *Xanthomonas translucens* pv. *translucens* is the causal agent of a bacterial wilt on *Hordeum vulgare* (barley) (1) that is often called black chaff. The strain DSM 18974^T (ATCC 19319^T) was isolated from its host plant in Minnesota, USA. It is the type strain of the species *X. translucens* (2). *X. translucens* strains occur worldwide and infect a broad range of cereals and forage grasses. At the same time, *Xanthomonas* strains are applied in biotechnology to synthesize the exopolysaccharide xanthan (3), a commercial thickening agent. Despite gradually rising numbers of publications, our knowledge about *X. translucens* genomes is still fragmentary. Hence, detailed genome data of *X. translucens* pv. *translucens* are suitable to advance functional research and deepen our phylogenetic understanding of *X. translucens* and related xanthomonads.

Initially, genomic DNA of *X. translucens* pv. *translucens* DSM 18974^T was extracted (4) to generate a paired-end library that was shotgun-sequenced with a Genome Sequencer FLX (GS FLX) system by means of the 454 Titanium technology (Roche) as described earlier (5). However, the assembly of the resulting data produced a large number of contigs, possibly pointing to the presence of repetitive IS elements. To address this deficiency, a combination of single-molecule real-time (SMRT) and Illumina sequencing technologies was applied. Genomic DNA was extracted applying Qiagen Genomic-tips 100/G according to the manufacturer's instructions and sequenced on a PacBio RSII instrument (Pacific Biosciences) using P5 chemistry. Genome assembly was performed with the "RS_HGAP_Assembly.3" protocol included in SMRT Portal version 2.3.0, utilizing 49,791 post-filtered reads with an average read length of 12,646 bp. One complete chromosomal contig was obtained and trimmed, circularized, and adjusted to the beginning of the *dnaA* gene. To improve the sequence quality, additional genome data were obtained by sequencing a

paired-end TruSeq PCR-free DNA fragment library with an average size of 719 ± 226 bp on a MiSeq instrument (Illumina) in a 2 × 300-bp run. This provided a total of 1,021,089,076 bp in 4,490,068 reads, among them 2,129,085 paired-ends reads. The Illumina reads were mapped with the Burrows–Wheeler aligner (6) onto the contig to obtain the consensus sequence of a circular chromosome of 4,715,357 bp with an average coverage of 223× and a G+C content of 67.7%. Automated genome annotation carried out by means of the GenDB platform (7) with the Prodigal gene prediction software (8) revealed 3,736 protein-coding genes, 54 tRNAs, and two rRNA operons.

Comparative analysis employing the EDGAR software (9) facilitated the identification of a CRISPR array in addition to virulence factor genes, which encoded protein-secretion systems like a *hrp* type III secretion system with transcription activator-like effectors (10) and the polysaccharide xanthan.

Accession number(s). This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession number [LT604072](https://www.ncbi.nlm.nih.gov/nuccore/LT604072). The version described in this paper is the first version, LT604072.1.

ACKNOWLEDGMENTS

We sincerely thank Simone Severitt and Nicole Heyer for technical assistance. The project benefited from the financial support of the German Federal Ministry of Education and Research, BMBF, for the project "Bielefeld-Gießen Center for Microbial Bioinformatics—BiGi" (grant 031A533) within the German Network for Bioinformatics Infrastructure (de.NBI). The funders had no role in study design, data collection, and interpretation, or in the decision to submit the work for publication. We acknowledge support for the article processing charge by the Deutsche Forschungsgemeinschaft and the Open Access Publication Fund of Bielefeld University.

FUNDING INFORMATION

This work, including the efforts of Daniel Wibberg, Anika Winkler, Alfred Pühler, and Frank-Jörg Vorhölter, was funded by Bundesministerium für Bildung und Forschung (Federal Ministry of Education and Research) (031A533).

REFERENCES

- Duveiller E, Maraite H. 1993. *Xanthomonas campestris* pathovars on cereals: cause of leaf streak or black chaff diseases, p. 76–79. In Swings JG, Civerolo EL (ed), *Xanthomonas*. Chapman and Hall, London, United Kingdom.
- Rademaker JL, Norman DJ, Forster RL, Louws FJ, Schultz MH, de Bruijn FJ. 2006. Classification and identification of *Xanthomonas translucens* isolates, including those pathogenic to ornamental asparagus. *Phytopathology* 96:876–884. <http://dx.doi.org/10.1094/PHTO-96-0876>.
- Becker A, Katzen F, Pühler A, Ielpi L. 1998. Xanthan gum biosynthesis and application: a biochemical/genetic perspective. *Appl Microbiol Biotechnol* 50:145–152. <http://dx.doi.org/10.1007/s002530051269>.
- Hopwood DA, Bibb MJ, Chater KF, Kieser T, Bruton CJ, Kieser HM, Lydiate DJ, Smith CP, Ward JM, Schrepf H. 1985. Genetic manipulation of streptomycetes—a laboratory manual. John Innes Foundation, Norwich, United Kingdom.
- Arnold M, Wibberg D, Blom J, Schatschneider S, Winkler A, Kutter Y, Rückert C, Albersmeier A, Albaum S, Goesmann A, Zange S, Heesemann J, Pühler A, Hogardt M, Vorhölter FJ. 2015. Draft genome sequence of *Pseudomonas aeruginosa* strain WS136, a highly cytotoxic ExoS-positive wound isolate recovered from pyoderma gangrenosum. *Genome Announc* 3(4):. <http://dx.doi.org/10.1128/genomeA.00680-15>.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics* 26:589–595. <http://dx.doi.org/10.1093/bioinformatics/btp698>.
- Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J, Kalinowski J, Linke B, Rupp O, Giegerich R, Pühler A. 2003. GenDB—an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res* 31:2187–2195. <http://dx.doi.org/10.1093/nar/gkg312>.
- Wichmann F, Vorhölter FJ, Hersemann L, Widmer F, Blom J, Niehaus K, Reinhard S, Conradin C, Kölliker R. 2013. The noncanonical type III secretion system of *Xanthomonas translucens* pv. *graminis* is essential for forage grass infection. *Mol Plant Pathol* 14:576–588. <http://dx.doi.org/10.1111/mpp.12030>.
- Blom J, Albaum SP, Doppmeier D, Pühler A, Vorhölter FJ, Zakrzewski M, Goesmann A. 2009. EDGAR: a software framework for the comparative analysis of prokaryotic genomes. *BMC Bioinformatics* 10:154. <http://dx.doi.org/10.1186/1471-2105-10-154>.
- Boch J, Bonas U, Lahaye T. 2014. TAL effectors—pathogen strategies and plant resistance engineering. *New Phytol* 204:823–832. <http://dx.doi.org/10.1111/nph.13015>.