

An mRNA Blueprint for C₄ Photosynthesis Derived from Comparative Transcriptomics of Closely Related C₃ and C₄ Species^{1[W][OA]}

Andrea Bräutigam², Kaisa Kajala², Julia Wullenweber, Manuel Sommer, David Gagneul, Katrin L. Weber, Kevin M. Carr, Udo Gowik, Janina Maß, Martin J. Lercher, Peter Westhoff, Julian M. Hibberd², and Andreas P.M. Weber^{2*}

Institute of Plant Biochemistry (A.B., J.W., M.S., D.G., K.L.W., A.P.M.W.), Institute of Plant Molecular and Developmental Biology (U.G., P.W.), and Institute of Informatics (J.M., M.J.L.), Heinrich-Heine University, 40225 Duesseldorf, Germany; Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, United Kingdom (K.K., J.M.H.); and Bioinformatics Core, Research Technology Support Facility, Michigan State University, East Lansing, Michigan 48824 (K.M.C.)

C₄ photosynthesis involves alterations to the biochemistry, cell biology, and development of leaves. Together, these modifications increase the efficiency of photosynthesis, and despite the apparent complexity of the pathway, it has evolved at least 45 times independently within the angiosperms. To provide insight into the extent to which gene expression is altered between C₃ and C₄ leaves, and to identify candidates associated with the C₄ pathway, we used massively parallel mRNA sequencing of closely related C₃ (*Cleome spinosa*) and C₄ (*Cleome gynandra*) species. Gene annotation was facilitated by the phylogenetic proximity of *Cleome* and *Arabidopsis* (*Arabidopsis thaliana*). Up to 603 transcripts differ in abundance between these C₃ and C₄ leaves. These include 17 transcription factors, putative transport proteins, as well as genes that in *Arabidopsis* are implicated in chloroplast movement and expansion, plasmodesmatal connectivity, and cell wall modification. These are all characteristics known to alter in a C₄ leaf but that previously had remained undefined at the molecular level. We also document large shifts in overall transcription profiles for selected functional classes. Our approach defines the extent to which transcript abundance in these C₃ and C₄ leaves differs, provides a blueprint for the NAD-malic enzyme C₄ pathway operating in a dicotyledon, and furthermore identifies potential regulators. We anticipate that comparative transcriptomics of closely related species will provide deep insight into the evolution of other complex traits.

C₄ photosynthesis is a complex biological trait that enables plants to either accumulate biomass at a much faster rate or live in adverse environments compared with “ordinary” plants (Hatch, 1987; Osborne and Freckleton, 2009). These C₄ plants have added a CO₂ concentration mechanism on top of their regular photosynthetic carbon fixation that makes them not only more efficient at assimilating inorganic carbon; they frequently also have higher water and nitrogen use efficiencies (Black, 1973; Oaks, 1994; Osborne and

Freckleton, 2009). Beyond the basic biochemistry, our understanding of C₄ photosynthesis is limited.

The principle of C₄ photosynthesis is deceptively simple: instead of using Rubisco as the primary carbon-fixing enzyme, C₄ plants use phosphoenolpyruvate carboxylase (PEPC). Unlike Rubisco, PEPC is more specific for inorganic carbon (Hatch, 1987). Since the C₄ cycle is an add-on rather than a replacement for Rubisco and the Calvin-Benson cycle, the prefixed CO₂ is transported in a bound form, a C₄ acid (hence the name), to the site of Rubisco. The C₄ cycle generates high concentrations of CO₂ around Rubisco (Hatch, 1987), and this increases the rate of photosynthesis because competition between CO₂ and oxygen at the active site of Rubisco is reduced (Jordan and Ogren, 1984). In most C₄ plants, concentrating CO₂ around Rubisco involves the reactions of photosynthesis being partitioned between bundle sheath (BS) and mesophyll (M) cells as well as changes to cell biology and leaf development (Hatch, 1987; Sage, 2004), although in some lineages, C₄ photosynthesis operates within individual cells (Reiskind et al., 1989; Keeley, 1998; Voznesenskaya et al., 2001, 2002, 2003).

In all known C₄ plants, CO₂ enters M cells and is converted into bicarbonate by carbonic anhydrase.

¹ This work was supported by the German Research Council (grant nos. WE 2231/4-1 to A.P.M.W., SFB TR1 to P.W. and A.P.M.W., and IRTG 1525/1 to P.W. and A.P.M.W.) and the Leverhulme Trust and Isaac Newton Trust (to J.M.H.).

² These authors contributed equally to the article.

* Corresponding author; e-mail andreas.weber@uni-duesseldorf.de.

The authors responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) are: Julian M. Hibberd (julian.hibberd@plantsci.cam.ac.uk) and Andreas P.M. Weber (andreas.weber@uni-duesseldorf.de).

[W] The online version of this article contains Web-only data.

[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.110.159442

PEPC then combines HCO₃⁻ with PEP to generate the C₄ oxaloacetic acid, which is rapidly converted into either Asp or malate. These C₄ acids then diffuse to the site of Rubisco through abundant plasmodesmata, where C₄ acid decarboxylases release CO₂ (Hatch, 1987). Three distinct C₄ acid decarboxylases, known as NADP-dependent malic enzyme (NADP-ME), NAD-dependent malic enzyme (NAD-ME), and PEP carboxykinase, have been coopted into the C₄ pathway, and this has been used to define three biochemical subtypes of C₄ photosynthesis. The three-carbon compound released after decarboxylation diffuses back to the M cells and is converted to PEP catalyzed by pyruvate, orthophosphate dikinase (PPDK; Hatch and Slack, 1968). Because the enzymes involved in the C₄ cycle are found in the cytosol, chloroplasts, and mitochondria, a significant amount of transport across organellar membranes is required for the C₄ cycle to operate. However, few genes encoding transporters that allow the increased intracellular flux of metabolites required for C₄ photosynthesis have been identified (Bräutigam et al., 2008a; Majeran and van Wijk, 2009). In addition, we have a very limited understanding of the mechanisms controlling the altered cell biology and morphology associated with C₄ leaves. The C₄ cycle likely affects not only the relatively small number of enzymes and transport proteins needed to perform the core reactions but, given the consequences on the ecological performance of the plants, also a range of other processes.

The gaps in our understanding of the mechanisms underlying C₄ photosynthesis limit insight into a metabolic pathway that has evolved repeatedly at least 45 times in plants (Sage, 2004) and so is of interest in terms of understanding a remarkable example of convergent evolution. In addition, because C₄ plants are among the most productive on the planet and the pathway is associated with increased water and nitrogen use efficiencies (Brown, 1999), it has been suggested that characteristics of C₄ photosynthesis should be placed into C₃ crops (Matsuoka et al., 2001; Mitchell and Sheehy, 2006; Hibberd et al., 2008). A more complete understanding of genes involved in C₄ photosynthesis is fundamental to attempts at placing components of the C₄ pathway into C₃ crops to increase yield.

Recently, a new set of tools has become available to analyze species without sequenced genomes on a genomic scale: next generation sequencing (NGS) technology (summarized in Metzker, 2010). With NGS, the transcriptome of a tissue can be sequenced and quantified at the same time (RNA-Seq; Wang et al., 2009). The 454 FLX genome sequencer provides a quarter million sequence reads of 230 bases in each run from a cDNA template generated from mRNA (<http://www.454.com/>; Metzker, 2010). The resulting reads can be mapped onto a closely related reference to quantify the number of reads matching a gene locus, thus providing a measure of transcript abundance (Flicek and Birney, 2009; Bräutigam and Gowik, 2010). We chose to compare the C₄ plant *Cleome gynandra* with the C₃ plant

Cleome spinosa, since they are members of the same genus and are closely related to *Arabidopsis* (*Arabidopsis thaliana*; Brown et al., 2005; Marshall et al., 2007). Given the close phylogenetic relationship, we can take advantage of the well-annotated *Arabidopsis* genome (Swarbreck et al., 2008) and its known genome history (Bowers et al., 2003; Haberer et al., 2004; Thomas et al., 2006) to identify and quantify the biological functions regulated at the level of transcript abundance in the C₄ species compared with the C₃ species. Although the experiment will also capture variation in the abundance of transcripts associated with differences between the species that do not relate to C₄ photosynthesis, the close proximity of the *Cleome* species should reduce this effect. We chose to use mature fully differentiated leaves for the analysis, since we wanted to minimize the influence of species-specific effects during leaf differentiation but rather focus on transcript profiles when C₄ photosynthesis is fully operational. Once this profile is defined, analysis of developmental stages may reveal how the profile is achieved during differentiation.

By comparing the transcriptomes of closely related C₃ and C₄ species, we will test (1) whether cross-species transcriptomic comparisons are feasible, (2) the degree to which the core C₄ cycle enzymes and transport proteins are regulated at the level of transcript abundance, and (3) whether the changes in metabolism associated with C₄ photosynthesis are associated with additional unexpected shifts in transcript profiles in leaves of C₄ compared with C₃ plants, and (4) define candidates for additional functions critical to C₄ photosynthesis based on unbiased observation of the data. By analyzing the complete transcriptome, we define the maximal extent to which the C₄ pathway alters leaf transcript profiles.

RESULTS

Physiological Analysis of C₃ and C₄ Leaves Confirms C₄ Metabolism in *C. gynandra*

To confirm that the *C. spinosa* and *C. gynandra* leaves we used for transcriptomic analysis were using C₃ and C₄ photosynthesis, respectively, we analyzed the steady-state levels of metabolites associated with the C₄ cycle. For example, large quantities of Asp, Ala, and pyruvate are produced in M and BS cells of NAD-ME C₄ leaves, and they were 19, 3.9, and 3.6 times more abundant, respectively, in *C. gynandra* compared with *C. spinosa* (Supplemental Table S1). In contrast, and in agreement with the lower demand for the photorespiration in C₄ leaves, glycerate and glycolate, intermediates of the photorespiratory cycle, were 4.5 and 1.9 times more abundant in *C. spinosa* (Supplemental Table S1). We also determined the extractable activities of PEPC, aspartate aminotransferase (AspAT), NAD-dependent malate dehydrogenase (NAD-MDH), NAD-ME, and alanine aminotransferase (AlaAT). Except for NAD-MDH, significantly higher activities of the enzymes required for the C₄ cycle were measured in

C. gynandra leaf extracts (Supplemental Fig. S1). The metabolite profiling of leaf extracts using gas chromatography-electron impact-time of flight (GC-EI-TOF) and the enzyme activity assays showed that the plants we used for digital gene expression analysis had clear differences in their metabolite profiles and enzyme activities, and these were consistent with functional C₃ and C₄ photosynthesis operating in leaves of *C. spinosa* and *C. gynandra*, respectively.

The Leaf Transcriptomes for Closely Related C₃ and C₄ Species Are Qualitatively Similar

To obtain sequence tags for digital gene expression (DGE) analysis from *C. spinosa* (C₃) and *C. gynandra* (C₄), RNA was isolated from mature leaves of each species and prepared for 454 sequencing. One sequencing run on a Genome Sequencer FLX (GS FLX; Roche) sequencing system was conducted on leaf cDNA isolated from either *C. gynandra* or *C. spinosa*. From *C. spinosa*, we obtained 70,564,592 nucleotides, and from *C. gynandra*, 91,851,136 nucleotides of raw sequence were obtained; after quality control, these corresponded to 65,525,139 and 85,681,233 nucleotides, respectively (Table I). The mean read length of the cleaned sequence reads was 232 nucleotides for *C. gynandra* and 230 nucleotides for *C. spinosa* (Table I).

To exclude program-specific mapping artifacts and to test whether the *C. gynandra* and *C. spinosa* libraries behave robustly during mapping, two different programs, BLAST and BLAT (BLAST-Like Alignment Tool), were used to align the reads to Arabidopsis as the reference genome. To define the most suitable mapping parameters, an array of parameters for mappings in both the DNA and protein space were tested (Table II). Neither the *C. gynandra* nor the *C. spinosa* library mapped well to Arabidopsis cDNAs in the DNA space using BLAT or BLAST, although the differences are more dramatic for BLAT (Table II). In the protein space, however, the proportion of mapped reads increased dramatically. When 75% amino acid

sequence identity was required, three-quarters of the reads could be mapped with BLAT, resulting in 1.48 and 1.57 average mappings per read, respectively. Even with the most lenient mapping parameters, the proportion of mapped reads did not exceed 83% with BLAT and 78.8% with BLAST (Table II). In all mapping attempts, the *C. gynandra* and *C. spinosa* read libraries yielded qualitatively similar mapping results, irrespective of mapping program or parameters.

To obtain a stringent yet inclusive mapping, the mapping conducted in protein space at 75% or greater identity with BLAT was chosen, and this mapping file was parsed by in-house scripts to keep only the read match with the highest number of matching bases. For a more lenient mapping, a BLAST mapping at a cutoff of $1e^{-5}$ was chosen and parsed to keep only the best BLAST hit for each read. For each Arabidopsis Genome Initiative (AGI) code, the number of matching reads was counted and the hit count was then transformed to reads per million (RPM) to normalize for the number of reads available for each species. After parsing, the sequenced libraries matched between 50.5% and 55.3% of the genes in the Arabidopsis reference (Supplemental Table S2).

To assess whether the data sets for the two different species and the two different mappings were qualitatively similar, we tested the coverage of the functional classes. Overall, about 50% of all genes were represented in both species with the BLAT (Fig. 1A) and the BLAST mapping (Fig. 1B). Although the majority of gene classes were represented by more than 50% of genes in each class for both mappings, the classes function unknown, putative lipid transfer protein, storage protein, and defense were underrepresented compared with all genes (Fig. 1). Genes present in the organellar genomes were not well represented (Supplemental Table S3). Genes classified into primary metabolism including photosynthesis, central carbon, nitrogen metabolism, amino acid, and nucleotide metabolism as well as many cellular processes were well-represented categories, and about four-fifths of genes predicted to be involved in the C₄ pathway were detected in both species. Overall, the pattern of detection in the different gene classes was similar for both species and independent of the program used for the mapping (Fig. 1).

Table I. Massively parallel signature sequencing allows large-scale assembly of transcripts in both *C. spinosa* and *C. gynandra* after comparison with the TAIR 8 Arabidopsis database

One GS FLX sequencing run allowed significant generation of sequence for both species, and the vast majority of these could be used to assemble contigs and then matched to Arabidopsis genes.

Data	<i>C. spinosa</i>	<i>C. gynandra</i>
Raw reads	313,807	402,674
Raw nucleotides	70,564,592	91,851,136
Raw mean length	225	228
Clean reads	284,318	368,333
Clean nucleotides	65,525,139	85,681,233
Clean mean length	230	232
Contigs	17,655	18,992
Total length (nucleotides)	7,746,894	9,062,043
Total reads	245,324	319,732
Percent assembled	86.3	86.8

Transcripts of Known C₄ Genes Are More Abundant with One Exception

Detailed analysis of known C₄ genes showed that all but one gene necessary for the core C₄ cycle of NAD-ME-type plants were massively up-regulated in *C. gynandra* compared with *C. spinosa*. Transcripts encoding PEPC were up-regulated 78-fold, those encoding AspAT were up-regulated 343-fold, the transcripts for the two isoforms of NAD-ME were up-regulated 27- and 21-fold, respectively, and AlaAT were up-regulated 29-fold (Table III). The results for the BLAT

Table II. Mapping the sequence reads with different BLAT and BLAST parameters to empirically determine suitable mapping conditions

The percentage of AGI codes with at least one mapped read and the average mappings per read were determined prior to parsing the tables to retain only the best match. Suitable mapping conditions are printed in bold; for BLAT, the cutoff value is the minimal number of matching bases; for BLAST, it is the minimal accepted e-value.

Mapping Program	Library	Search Space	Cutoff Value	Percentage Reads with at Least One Hit in the Reference	Percentage AGI Codes with at Least One Mapped Read	Average Mappings per Read
BLAT	<i>C. gynandra</i>	DNA	60	40.9	42.0	1.19
			75	40.7	41.7	1.19
			85	30.2	35.8	1.15
			90	7.7	19.5	1.09
		Protein	25	82.6	70.4	2.35
			50	82.6	70.4	2.35
			75	75.4	62.6	1.48
			80	56.4	52.2	1.27
	<i>C. spinosa</i>	DNA	60	40.8	38.9	1.29
			75	40.6	38.5	1.28
			85	29.7	32.3	1.21
			90	8.5	17.1	1.15
		Protein	25	83.0	67.7	2.49
			50	83.0	67.7	2.46
			75	76.0	58.9	1.57
			80	57.9	48.4	1.32
BLAST	<i>C. gynandra</i>	DNA	1e-05:	68.9	56.5	30.7
			1e-10:	58.8	49.1	27.7
			1e-30:	29.6	30.5	18.9
			1e-50:	9.9	15.9	11.5
		Protein	1e-05:	78.0	76.9	106.6
			1e-10:	67.8	71.0	64.6
			1e-30:	29.0	39.5	22.9
			1e-50:	0.1	0.3	7.5
	<i>C. spinosa</i>	DNA	1e-05:	69.7	53.0	28.2
			1e-10:	59.6	46.3	25.1
			1e-30:	29.4	28.3	16.2
			1e-50:	9.8	14.4	9.8
		Protein	1e-05:	78.8	75.3	93.7
			1e-10:	68.3	68.7	56.4
			1e-30:	29.3	36.0	21.2
			1e-50:	0.1	0.3	4.6

and the BLAST mappings were similar with one exception. In the BLAST mapping, the reads mapping to PEPC were split onto two genes in the Arabidopsis reference genome, whereas they mapped to only one gene in the BLAT mapping (Table III). Transcripts encoding mitochondrial malate dehydrogenases were increased only 1.3-fold (Supplemental Table S3). Not only were genes associated with the C₄ pathway up-regulated compared with C₃, but they also had high absolute read counts between 1,800 and 4,806 RPM.

The Leaf Transcriptomes for Closely Related C₃ and C₄ Species Are Quantitatively Different

Before undertaking detailed analysis of differences in transcript abundance between *C. gynandra* and *C. spinosa*, we used quantitative (q)PCR to confirm estimates of transcript abundance identified by RNA-Seq.

We chose genes whose transcript abundance differed over 4 orders of magnitude and used qPCR to assess their abundance. qPCR was performed on both the cDNA used for RNA-Seq and cDNA generated from RNA isolated from leaves in a separate experiment. This approach provided strong support for the differences in abundance of transcripts between the two species that we determined from RNA-Seq (Fig. 2). Overall, this showed that the ratios of transcript abundance obtained by RNA-Seq-based DGE are suitable for calling differentially expressed genes between two related species.

Of the 13,662 transcripts for which we captured quantitative data (Supplemental Table S3), we identified 583 (BLAT) or 603 (BLAST) transcripts whose abundance differed significantly ($P \leq 0.01$) between *C. spinosa* and *C. gynandra*, with 256/258 (1.2%/1.2%) transcripts being more abundant in *C. gynandra* and




Figure 1. The qualitative patterns of transcript abundance between *C. gynandra* and *C. spinosa* are very similar, with the same classes underrepresented and overrepresented in both libraries. A, Analysis based on BLAT mapping. B, Analysis based on BLAST mapping. Black bars refer to the C_4 plant *C. gynandra*, and white bars refer to the C_3 plant *C. spinosa*.

327/345 (1.5%/1.6%) transcripts being more abundant in *C. spinosa* (Fig. 3, "all"). We tested whether significantly changed transcripts are enriched in functional categories and whether they were more highly expressed in the C_4 or the C_3 species. While the qualitative classification of detected genes showed a very similar pattern between *C. spinosa* and *C. gynandra* (Fig. 1), the quantitative analysis revealed massive differences in representation between gene classes in the C_3 and the C_4 species (Fig. 3). The transcript profile generated by the BLAT mapping (Fig. 3A) is similar to the one generated by the BLAST mapping (Fig. 3B), although not all genes called as significantly regulated were identical (Supplemental Table S3). The classes containing the highest percentage of changed genes are the photosynthetic classes as well as the C_4 cycle, Calvin-Benson cycle, and photorespiration (Fig. 3). The latter two have lower steady-state mRNA levels in C_4 leaf tissue (Fig. 3, bottom), while the photosynthetic

classes of PSI, cyclic electron flow, and cytochrome b_6/f complex as well as the C_4 cycle have higher levels in C_4 leaf tissue (Fig. 3, top). A number of classes involved in primary metabolism also have lower steady-state transcript levels in C_4 tissues: one-carbon compound metabolism, other central carbon metabolism, shikimate pathway, and amino acid metabolism. Protein synthesis also has lower steady-state transcript levels, which are limited to cytosolic and plastidic protein synthesis genes (Supplemental Fig. S3). Among the classes with higher steady-state transcript levels are starch metabolism, cofactor synthesis, putative lipid transfer proteins, nitrogen metabolism, and β -1,3 glucan metabolism. The quantitative pattern (Fig. 3) is similar to the qualitative pattern (Fig. 1) with regard to the influence of the mapping program; the BLAT and BLAST mappings look remarkably similar with the exception of shikimate metabolism.

Table III. Transcript abundance of C₄ cycle genes that have significantly higher transcript abundance in C₄ leaf tissue

Asterisks denote changes significant only in BLAST mapping.

Enzyme	Locus	BLAT Mapping			BLAST Mapping		
		<i>C. gynandra</i> RPM	<i>C. spinosa</i> RPM	Fold Change	<i>C. gynandra</i> RPM	<i>C. spinosa</i> RPM	Fold Change
AspAT	AT2G30970	4,806	14	343.3	4,601	18	257.9
PPDK	AT4G15530	3,262	14	233.0	3,216	13	240.3
PEPC	AT2G42600	9,702	124	78.2	8,321	169	49.1
AlaAT	AT1G17290	7,610	267	28.5	7,242	259	28.0
NAD-ME1	AT4G00570	1,357	51	26.6	1,326	49	27.0
NAD-ME2	AT2G13560	1,800	87	20.7	1,723	85	20.3
PEPC kinase	AT1G08650	230	37	6.2	226	36	6.3
NADP-ME*	AT1G79750	227	60	3.8	216	45	4.8
PEPC*	AT1G53310	94	248	0.4	950	192	5.0
PPDK regulatory protein*	AT4g21210	148	32	4.6	198	27	7.3

Transcripts with Similar Patterns of Abundance Compared with Bona Fide C₄ Genes and Rubisco

The list of 13,662 transcripts detected in either *C. spinosa* or *C. gynandra* tissues and the list of 603 transcripts that are differentially regulated between both species (Supplemental Table S3, BLAST mapping) prompted us to determine which transcripts showed changes in abundance similar to the core C₄ genes or Rubisco subunit-encoding genes. Such transcripts display both a large fold change between the C₃ and the C₄ plants and large absolute read numbers. For example, among the transcripts encoding putative transport proteins, three plastidic transport proteins, the PEP phosphate translocator PPT, a putative bile acid:sodium symporter, and a putative proton:sodium antiporter, two mitochondrial dicarboxylate carriers, and one plasma membrane intrinsic protein were massively up-regulated in C₄ *C. gynandra* (Table IV). No transcripts encoding transport proteins were found to be down-regulated to a comparable degree. Among metabolic genes, two cytosolic carbonic anhydrases, one of which (CA4; Table IV) is likely tethered to the plasma membrane, an adenylate kinase, and a pyrophosphatase were up-regulated at levels comparable to those of C₄ cycle genes. Many proteins of unknown function showed differential expression, the most striking case being a putative lipid transfer protein, also annotated as an extensin-like protein. Based on annotation and differential expression pattern, several transcripts predicted to encode known C₄ functions that have not yet been assigned to genes, such as CHLOROPLAST UNUSUAL POSITIONING1 (CHUP1) and actin for chloroplast positioning or callose-degrading enzymes for regulating plasmodesmatal opening, were identified (Table IV).

Regulatory Genes That Are Significantly Changed

The transcript profiles of these C₃ and C₄ species identify a number of regulatory proteins that are candidates for maintaining C₄ status. Among transcripts encoding proteins with regulatory functions, 43

were significantly up-regulated in either *C. gynandra* or *C. spinosa* (Fig. 3). These include bona fide transcription factors, protein phosphatases and kinases, and the regulatory proteins of the pyruvate dehydrogenase complex (up-regulated in C₄), of PPDK (up-regulated in C₄), and of Rubisco (down-regulated in C₄). Only 17 transcription factors are significantly changed; seven of those have higher steady-state mRNA levels compared with the C₃ leaf tissue, while 10 have lower steady-state mRNA levels (Table V).

In addition to the detailed quantitative and qualitative analysis of read mappings to generate ESTs for both species, contigs were assembled from cleaned reads for each species as described previously (Weber et al., 2007; Bräutigam et al., 2008b) and then annotated by BLASTX versus The Arabidopsis Information Resource (TAIR) 9 protein models. A total of 18,992 and 17,655 contigs representing total sequence lengths of 9,062,043 and 7,746,894 nucleotides were obtained for *C. gynandra* and *C. spinosa*, respectively (Table I).




Figure 2. Massively parallel sequencing of mRNAs (RNA-Seq) and qPCR generate similar profiles of transcript abundance in *C. gynandra* and *C. spinosa*. Ratios of transcript abundance in *C. gynandra* and *C. spinosa* were calculated, and transcripts selected for this analysis spanned 4 orders of magnitude. CA, Carbonic anhydrase; PPCK, PEPC kinase; LHCA, light-harvesting complex subunit A; RbcS1a, ribulose biphosphate carboxylase oxygenase 1a; RCA, Rubisco activase. Black bars represent data from RNA-Seq, and white bars represent data from qPCR. The horizontal dashed line represents a ratio of 1 and indicates no difference in transcript abundance between the two species.




Figure 3. The quantitative patterns of transcript accumulation in *C. gynandra* and *C. spinosa* are distinct. A, Analysis based on BLAT mapping. B, Analysis based on BLAST mapping. Shown are the percentages of genes with significantly higher abundance of transcripts in C_4 (red bars), unchanged (white bars, including genes not detected), and significantly lower abundance of transcripts in C_4 (blue bars) based on the total number of genes in each annotation class (in parentheses on the y axis).

DISCUSSION

Transcriptomic Comparisons of Different Species with NGS Technology Are Feasible

Read mapping by alignment is a well-established tool to quantify transcript abundance and thus determine mRNA steady-state levels (Wall et al., 2009; Metzker, 2010). The concept of mapping to a cross-species reference has also been established theoretically (Palmieri and Schlotterer, 2009), although the potential has not been experimentally explored to date (Bräutigam and Gowik, 2010).

To explore cross-species mapping, the transcriptome sequencing was carried out using 454 FLX, a long-read technology, since theoretical work had established that at least BLAT is capable of mapping reads that contain alterations in comparison with the reference if the reads are at least 100 bases long (Palmieri and Schlotterer,

2009). We also established a reference database, which removes the genome history of *Arabidopsis* as far as it is known (Bowers et al., 2003; Haberer et al., 2004; Thomas et al., 2006). Tandem duplicated genes and segmentally duplicated genes (remnants of the last whole genome duplications) were removed to prevent genome history from interfering with comparative quantitative mapping (Bräutigam and Gowik, 2010).

Both BLAT and BLAST mappings indicate that using a minimal reference does not diminish read mappings (Supplemental Table S4) while avoiding mapping problems based on genome history (Bräutigam and Gowik, 2010). The mappings in protein space allowed more successful read mappings, because protein sequences diverge more slowly than nucleotide sequences. Although the proportion of reads mapped varied with changing mapping parameters (Table II;

Table IV. Transcript abundance of selected genes with an expression similar to that of C₄ cycle genes and RubiscoAll changes are significant at $P \leq 0.01$. n/a, Not available.

Function	Locus	Annotation (TAIR 9)	<i>C. gynandra</i> RPM	<i>C. spinosa</i> RPM	Ratio
Transport proteins					
	AT2G26900	Bile acid:sodium symporter family protein	4,774	55	86.8
	AT2G22500	Mitochondrial dicarboxylate carrier	324	0	n/a
	AT4G24570	Mitochondrial dicarboxylate carrier	148	0	n/a
	AT2G45960	Plasma membrane intrinsic protein subfamily protein	2,686	133	20.2
	AT5G33320	Phosphoenolpyruvate/phosphate translocator	1,955	97	20.2
	AT1G49810	Member of Na ⁺ /H ⁺ antiporter family	1,321	83	15.9
Metabolism					
	AT3G52720	α -Carbonic anhydrase 1	227	152	1.5
	AT1G23730	β -Carbonic anhydrase 4	497	87	5.7
	AT5G35170	Adenylate kinase family protein	1,994	235	8.5
	AT5G09650	Inorganic pyrophosphatase	2,664	833	3.2
Proteins of unknown function					
	AT1G12090	Extensin-like protein (ELP)	6,278	147	42.7
Callose-degrading enzymes					
	AT3G57240	Member of glycosyl hydrolase family 17, likely β -1,3 glucanase	436	0	n/a
	AT1G32860	Member of glycosyl hydrolase family 17, likely β -1,3 glucanase	50	0	n/a
	AT5G42100	Plasmodesmal-associated β -1,3-glucanase	173	32	5.4
Cell biology					
	AT3G25690	CHUP1	22	170	0.13
	AT3G12110	ACTIN	122	727	0.2

Supplemental Table S4), the *C. spinosa* and *C. gynandra* libraries yielded similar results, indicating that, evolutionarily, both species are approximately equally distant from Arabidopsis, with mapping incurring similar penalties depending on parameters.

Since no read alignment program has emerged as the consensus program for NGS data analysis, two different programs were used for mapping and the output was compared in all cases. The output proved robust against changing the mapping program both qualitatively and quantitatively. When we mapped the quarter million reads obtained from each species of *Cleome* to a minimized TAIR 9 release of the Arabidopsis genome, they corresponded to approximately 11,000 loci. As the minimized TAIR 9 data set contains 21,972 gene loci, the reads we collected in *C. gynandra* and *C. spinosa* represent approximately 50% of the transcriptome. In Arabidopsis seedlings, approximately 60% of the loci represented in the TAIR 8 release were detectable (Weber et al., 2007); hence, we have likely captured a large proportion of the transcripts associated with leaves of *C. spinosa* and *C. gynandra*.

The qualitative representation of gene classes detected reflects that leaf tissues were analyzed. While photosynthetic genes as well as primary metabolism are well represented in all data sets, genes implicated in cell walls, secondary metabolism, and defense responses are underrepresented (Fig. 1). These classes contain genes that are likely specific to certain tissues, developmental stages, or environmental challenges. For example, cell wall genes may be better represented if our sampling had included expanding leaf or stem material (Schmid et al., 2005), and stress-response genes may be better represented if plants were sampled after exposure to extreme conditions (Kilian et al., 2007).

Likewise, certain pathways of secondary metabolism are likely restricted to defined tissues or developmental stages, making it unlikely that we would pick up many of these genes when profiling leaf libraries. Based on the gene detection pattern, the two plant species did not encounter different biotic or abiotic stresses or were not in different stages of growth, as very similar genes were detected in both species (Figs. 1 and 3).

Finally, only a very small proportion of transcripts showed significant differences in abundance between the two different species (Supplemental Tables S2 and S3), and these changes were enriched in a limited number of functional classes (Fig. 3). We conclude that cross-species mapping in protein space is a feasible strategy to compare different species as long as an equidistant reference is available.

Transcripts Derived from Core C₄ Cycle Genes Are More Abundant in the C₄ Species

C₄ photosynthesis has evolved convergently in many different lineages of plants (Sage, 2004), and in many cases the alterations to expression of specific genes has been related to transcriptional regulation (summarized in Sheen, 1999). Our genome-scale analysis allowed us to compare the steady-state transcript levels for all candidate C₄ genes at the same time. For all of the enzymes where a change in total extractable activity could be shown (Supplemental Fig. S1), a higher mRNA level of at least one isoform as judged from the read count was also present (Table III). The only enzyme showing no changes in transcript level is the mitochondrial NAD-MDH. Possibly, the activity of the mitochondrial NAD-MDH is high enough already

Table V. Transcription factors that are significantly changed between the leaf tissue samples

Asterisks denote changes significant only in BLAST mapping. n/a, Not available.

Locus	Transcription Factor Type	BLAT Mapping			BLAST Mapping			Segmentally Duplicated?
		<i>C. gynandra</i> RPM	<i>C. spinosa</i> RPM	Ratio	<i>C. gynandra</i> RPM	<i>C. spinosa</i> RPM	Ratio	
AT1G25560	AP2-EREBP	176	9	19.6	219	9	24.3	Yes
AT5G07580	AP2-EREBP	223	51	4.4	292	36	8.1	Yes
AT1G53910	AP2-EREBP*	32	138	0.2	84	268	0.3	Yes
AT5G10570	bHLH	0	83	n/a	0	112	n/a	Yes
AT3G21330	bHLH*	0	74	n/a	0	107	n/a	
AT3G62420	bZIP	11	138	0.1	10	138	0.1	
AT2G20570	G2-like	220	0	n/a	292	0	n/a	
AT1G72030	GNAT	11	179	0.1	10	330	0.0	
AT2G22430	HB	515	106	4.9	505	116	4.4	Yes
AT1G10200	LIM	22	230	0.1	21	205	0.1	
AT4G30410	Not specified*	0	32	n/a	0	76	n/a	
AT1G32700	PLATZ	176	9	19.6	115	4	28.8	
AT5G02810	Pseudo ARR-B	0	106	n/a	10	112	0.1	
AT2G36990	Sigma70-like	130	0	n/a	143	0	n/a	
AT1G48500	Tify	11	147	0.1	10	174	0.1	Yes
AT1G17380	Tify*	18	110	0.2	24	161	0.1	Yes
AT3G02790	Zinc finger	374	87	4.3	407	112	3.6	Yes

in C_3 plants to support a C_4 -type metabolic flux. The only transport protein known to date that is involved in the C_4 cycle, the PEP phosphate translocator (Fischer et al., 1997; Bräutigam et al., 2008a), is also up-regulated 20-fold, indicating that this transport protein is regulated at the level of mRNA abundance. Based on similarities in transcript abundance to known C_4 genes, our comparative RNA-Seq also identified likely additional components needed for C_4 photosynthesis. When PPK was characterized, it was proposed that adenylate kinase as well as inorganic pyrophosphatase need to be abundant in C_4 chloroplasts (Hatch and Slack, 1968). RNA-Seq confirmed this prediction and showed that the up-regulation also occurs at the level of transcript abundance. Taken together, we found that almost all transcripts encoding the proteins required for the core C_4 cycle have higher steady-state mRNA levels, and we propose that, at least in *C. gynandra*, the activity of C_4 cycle enzymes and transport proteins is controlled at least partially at the level of transcript abundance.

Alterations to the Abundance of Transcripts Associated with Other Metabolic Processes

Changes in the abundance of transcripts that are not associated with the core C_4 cycle are also detectable in leaves of *C. gynandra* and *C. spinosa*. The high-flux C_4 cycle poses additional demands in terms of ATP and reduction equivalents on the light reaction (Hatch, 1987). Specifically, the recycling of the initial CO_2 acceptor PEP requires additional ATP molecules (Hatch, 1987). In C_4 leaf tissue, one-third to one-half of the genes in the photosynthetic gene classes that contribute to ATP production by cyclic electron flow are up-regulated compared with C_3 leaf tissue: PSI, the

cytochrome b_6/f complex, and the genes mediating cyclic electron flow themselves (Fig. 3). It remains an open question whether these higher steady-state levels are caused by higher ATP demand or whether C_4 photosynthesis requires up-regulation of these genes to meet the ATP demand prior to establishing C_4 photosynthesis.

On the other hand, the classes of Calvin-Benson cycle genes and photorespiratory genes are those with the highest number of genes with significantly lower steady-state mRNA levels. It is a well-established fact that most C_4 plants have less Rubisco protein compared with C_3 plants (Ku et al., 1979) and that flux through the photorespiratory pathway is reduced compared with C_3 species (Chollet and Ogren, 1975; Leegood, 2002). Transcripts encoding the large and small subunits of Rubisco were reduced from 22,968 and 15,442 RPM to 6,984 and 4,900 RPM in *C. spinosa* and *C. gynandra*, respectively. Overall, the trend for Calvin-Benson cycle genes was for them to be down-regulated in *C. gynandra* compared with *C. spinosa* (Fig. 3). Likewise, a large number of genes encoding photorespiratory proteins, proteins involved in one-carbon compound metabolism, and the genes involved in ammonia re-assimilation, Gln synthetase, and Glu synthase have lower steady-state transcriptional levels (Fig. 3; Supplemental Table S3). The reduced flow through the photorespiratory pathway obviously decreases the demand on the expression system to maintain high steady-state levels of mRNA for many Calvin-Benson cycle and photorespiratory genes. The photosynthetic genes, the Calvin-Benson cycle and photorespiratory genes (in C_3), and the C_4 cycle genes (in C_4) are those with the highest read counts of the genes with known function (Supplemental Table S3). Although it is currently not possible to quantify

absolute transcript levels, since the genome of neither *Cleome* species has been sequenced, the high read counts obtained for the genes of central carbon metabolism and photosynthesis indicate that the steady-state levels of transcripts are high. Since the most altered gene classes are also those that contain the genes with the highest absolute read counts, it is not clear whether C₄ photosynthesis lowers or raises the demand on protein synthesis and accessory pathways such as amino acid synthesis. However, both the protein synthesis and the amino acid metabolism classes contain more genes that have lower steady-state levels in C₄ leaf tissue (Fig. 3). Within the protein synthesis gene class, many transcripts encoding structural components of plastidic and cytosolic ribosomes were reduced (Supplemental Fig. S3). This was not the case for components of mitochondrial ribosomes (Supplemental Fig. S3), indicating that there is not a general effect on translation but that the effect is likely specific to ribosomes involved in translation for the Calvin-Benson cycle and photorespiration. The protein-to-fresh weight ratio is also lower in C₄ leaf tissue compared with C₃ leaf tissue (Supplemental Fig. S2). We propose that plastidic ribosomes are relieved of the high translation load associated with the large subunit of Rubisco and that the cytosolic ribosomes need to translate fewer transcripts associated with central carbon metabolism as well as the small subunit of Rubisco. The reduced production of proteins in the leaves of C₄ plants is considered important in increasing nitrogen use efficiency, because the rate of photosynthesis per unit of nitrogen in the leaf is increased (Oaks, 1994). Our data indicate that there is also likely a significant saving in the nitrogen provision in the leaf, because fewer ribosomes as well as fewer proteins for central carbon metabolism are required.

The data set contains two additional gene classes, β -1,3 glucan metabolism and putative lipid transfer proteins, that showed differences in transcript abundance between *C. gynandra* and *C. spinosa* that could be explained within the current framework of knowledge of C₄ photosynthesis. The C₄ pathway requires efficient exchange of metabolites between M and BS cells via large numbers of plasmodesmata connecting both cell types, while the BS cell wall of many C₄ plants is suberized to reduce diffusion of CO₂ away from Rubisco (Hatch, 1987). Transcripts encoding three distinct glucan 1,3- β -glucosidases (Table IV) involved in governing plasmodesmatal conductivity by regulating the turnover of the β -1,3-glucan callose (Levy et al., 2007) were up-regulated in leaves of *C. gynandra* compared with *C. spinosa*. Therefore, it is possible that these genes are involved in increasing the open probability of plasmodesmata (Roberts and Oparka, 2003), which allows the efficient flux of organic acids between M and BS cells required during C₄ photosynthesis (Evert et al., 1977; Botha, 1992; Roberts and Oparka, 2003). A transcript annotated as a putative lipid transfer protein is among those that are most highly up-regulated in *C. gynandra* compared with *C.*

spinosa. Lipid transfer proteins are required for the export of lipids to the cell wall during cutin biosynthesis (DeBono et al., 2009). Interestingly, in *Arabidopsis*, some lipid transfer proteins are exclusively and abundantly expressed in the root endodermis, where suberin biosynthesis is required to establish the Casparian strip.

There are additional changes in the transcript profile that are less easily explained. Among the gene classes containing more genes with significantly higher transcript levels in C₄ leaf tissue are starch metabolism, cofactor synthesis and nitrogen metabolism, and heat shock/protein folding (in order of decreasing number of significantly different genes). On the other hand, it is difficult to conceive why genes involved in metal handling are frequently lower in transcript level in C₄ leaf tissues (Fig. 3). These changes may be connected to currently unknown phenomena relating to the C₄ pathway or may be part of differences not relating to C₄ photosynthesis between the two species. Overall, the global analysis of transcription on the level of functional classes reveals unexpected shifts in transcript profiles that can be explained based on the current knowledge about the C₄ pathway, while a range of smaller changes remain enigmatic.

Finally, our global transcriptional analysis of C₄ and C₃ leaf tissues not only allows testing hypotheses about the C₄ pathway on a global scale but also allows genes with expression patterns similar to those of known C₄ genes to be identified. The phylogenetic proximity of the Cleomaceae to *Arabidopsis* allows the identification of the orthologs in *Arabidopsis*, which will facilitate translational research into the model species (Brown et al., 2005).

Candidates for Additional C₄-Related Genes

The identification of transport proteins involved in the C₄ cycle lags behind that of enzymes, considering that the C₄ cycle requires the intracellular transport of pyruvate, PEP, Asp, and Ala across different organellar membranes (Bräutigam and Weber, 2011). A wide range of C₄ plants take up pyruvate into chloroplasts from the M in cotransport with sodium (Aoki et al., 1994; Aoki and Kanai, 1997), which might explain the requirement for sodium as a micronutrient in many C₄ species (Brownell and Crossland, 1972). Since the rate of pyruvate transport into C₄ M cell chloroplasts occurs at or exceeds the apparent rate of CO₂ assimilation, sodium-coupled pyruvate import implies a large influx of sodium into these chloroplasts, but the transporter has not yet been identified at the molecular level (Aoki and Kanai, 1997). Our finding that a putative plastidic proton:sodium symporter (NHD1) is 16-fold up-regulated in *C. gynandra* prompts us to hypothesize that it functions in exporting sodium from the chloroplast in order to maintain the sodium gradient required for import of pyruvate. In addition, we found strong up-regulation of a putative bile acid: sodium cotransporter in *C. gynandra*. Interestingly, up-

Table VI. Comparison of alterations in transcript abundance in C_4 and C_3 leaves with those induced by cold, sugar feeding, attack by pests or pathogens, diurnal changes to light, or circadian rhythms

Cause	Estimated Change in Transcriptome	Change	Reference
		%	
Cold treatment	514 (24,000) ATH1	2.1	Vogel et al. (2005)
C_4 leaves and C_3 leaves	583/603 (13,443/13,662)	2.7/2.8	This study
Glc feeding	978 (22,500) ATH1	4.4	Price et al. (2004)
<i>Pseudomonas syringae</i>	2,034 (23,750) ATH1	8.6	De Vos et al. (2005)
<i>Myzus persicae</i>	2,181(23,750) ATH1	9.1	De Vos et al. (2005)
Diurnal regulation	1,115 (11,521) cDNA array	11	Schaffer et al. (2001)
Circadian regulation	2,282 (18,890) Galbraith	12	Dodd et al. (2007)

regulation of the putative bile acid:sodium cotransporter or of NHD1 was not observed in maize (*Zea mays*; Bräutigam et al., 2008a), which belongs to a group of C_4 plants that show proton-dependent, not sodium-dependent, transport of pyruvate into M cell chloroplasts (Aoki et al., 1994; Aoki and Kanai, 1997). PEP generated from pyruvate in M cell chloroplasts is exported from these chloroplasts by PPT, thereby providing the substrate for the cytosolic PEPC reaction. Accordingly, transcripts encoding PPT are 20-fold up-regulated in *C. gynandra*, likely reflecting the increased requirement for transport of PEP (Table III). In contrast to what has been observed for the NADP-ME-type C_4 plant maize by quantitative proteomic analysis (Bräutigam et al., 2008a), we did not detect increased transcript abundance of the putative M chloroplast oxaloacetate/malate exchanger DiT1 (Taniguchi et al., 2002, 2004; Renne et al., 2003; Supplemental Table S3). This is consistent with the fact that oxaloacetic acid/malate shuttling across the M cell chloroplast envelope membrane is not required for NAD-ME-type C_4 photosynthesis (Weber and von Caemmerer, 2010; Bräutigam and Weber, 2011). The mitochondrial dicarboxylate carriers are prime suspects for the C_4 acid importer into the mitochondria, where decarboxylation takes place (Table IV). The initial uptake of inorganic carbon and its conversion to bicarbonate may be facilitated by the concerted action of a membrane intrinsic protein channeling the gas and a carbonic anhydrase that is predicted to be membrane bound (Table IV).

Chloroplasts in the BS of *C. gynandra are larger than those in the BS of C_3 species and, as in many other C_4 plants, are positioned in a strictly centripetal pattern (Marshall et al., 2007; Voznesenskaya et al., 2007). Transcripts derived from the *GIANT CHLOROPLAST1* (*GC1*) gene were more abundant in *C. gynandra* than in *C. spinosa* (Table IV). Although overexpression of *GC1* in Arabidopsis is reported not to effect chloroplast division (Maple et al., 2004), it is possible that it does so in *C. gynandra*. In addition, we also detected reduced accumulation of transcripts derived from the *CHUP1* and *ACTIN11* genes. In Arabidopsis, the outer chloroplast envelope membrane protein CHUP1 contains an actin-binding motif and is required for preventing chloroplast aggregation (Oikawa et al., 2003). Differential positioning of chloroplasts in BS and M*

cells of the C_4 plants finger millet (*Eleusine coracana*) and maize requires the actomyosin system (Kobayashi et al., 2009). Since AtCHUP1 is involved in positioning chloroplasts at the periclinal plasma membrane during the weak-light acclimation response via a coiled-coil domain and interaction with the cytoskeleton (Oikawa et al., 2003), it is possible that the centripetal positioning of chloroplasts in BS cells is linked to lower expression of the *CgCHUP1* and *ACTIN11* genes.

Controlling and Maintaining a C_4 State in Leaf Tissue

Our estimate that around 603 transcripts accumulate differentially in leaves of C_3 and C_4 species provides insight into the extent to which gene expression profiles change in C_4 leaves. For example, the fact that 258 transcripts were more abundant in the leaves of C_4 compared with C_3 species indicates that about 2.8% of the leaf transcriptome differentially accumulates in C_4 leaves (Supplemental Tables S2 and S6). To compare the complexity of the C_4 pathway with other multi-genic traits, we assessed the number of transcripts that are known to be regulated by sugars, cold, diurnal and




Figure 4. Schematic of components associated with the C_4 cycle in the NAD-ME subtype based on interpretation of RNA-Seq. Proteins that have been described previously are in gray, and novel proteins are marked in red. Metabolites are in black. PIP1B:CA4, PIP1B plasma membrane aquaporin:membrane-tethered carbonic anhydrase; OAA, oxaloacetic acid; ACT11-CHUP11, ACTIN11-CHUP1 complex; Pyr, pyruvate; OEP24, chloroplast outer envelope protein 24.

circadian rhythms, as well as attack by pests and pathogens (Table VI). Interestingly, the alterations in transcript abundance of leaves of *C. gynandra* compared with those of *C. spinosa* were greater than those observed in response to cold treatment and lower than those induced by Glc feeding, those occurring during pathogen attack, and the response to both diurnal and circadian rhythms. As significant progress has been made in understanding sugar signaling (Rolland et al., 2006), pathogen attack (Wise et al., 2007), and the control of gene expression in response to the diurnal cycle and circadian rhythms (Imaizumi et al., 2007), it should be possible to identify the regulators responsible for these alterations in transcript abundance in a C₄ leaf compared with a C₃ leaf. The changes in transcript abundance that we document in a C₄ leaf compared with a C₃ leaf likely overrepresent the changes in transcript abundance actually associated with C₄ photosynthesis on a whole leaf basis, as some differences in gene expression are likely due to the phylogenetic distance between *C. gynandra* and *C. spinosa*. A more confident estimate of the extent to which the leaf transcriptome is altered in association with C₄ photosynthesis will be generated when additional congeneric pairs of C₃ and C₄ species are subjected to deep transcriptome analysis and shared transcripts are identified. Between M and BS cells, the alterations in gene expression may be greater than those that we have defined for whole leaves. For example, up to 18% of genes are estimated to be differentially expressed between M and BS cells of maize (Sawers et al., 2007). However, it is not clear how different the transcript profiles of M and BS cells are in a dicot C₃ leaf, and until this is defined, it is not possible to infer the extent to which transcript abundance alters in these cell types in association with C₄ photosynthesis.

As we sampled from mature leaves to capture the differences between C₃ and C₄ leaves at the point of fully differentiated pathways, we likely also captured regulatory genes needed to maintain C₄ architecture and metabolism in mature leaves. Of the 17 transcription factors significantly altered (Table V), *GOLDEN2-LIKE1* (*GLK1*) has previously been implicated in regulating genes important in C₄ photosynthesis. In maize, *GOLDEN2* controls functional differentiation of chloroplasts in BS cells (Langdale and Kidner, 1994), and *GLK1* has been implicated in the expression of photosynthesis genes in M cells (Rossini et al., 2001). The fact that *GLK1* transcripts are significantly more abundant in leaves of *C. gynandra* would not necessarily be predicted, as previous work indicates that it becomes specialized in BS cells of C₄ leaves but not that its abundance is altered significantly. This implies that the increase in abundance of *GLK1* transcripts may not simply be due to its involvement in C₄ photosynthesis. When overexpression of *GLK1* was induced in *Arabidopsis*, the abundance of 114 transcripts was altered (Waters et al., 2009). We assessed the extent to which the genes that are controlled by *GLK1* change in abundance in leaves of *C. gynandra*

compared with *C. spinosa* and found that only 19 genes were shared between the two data sets. This may be due to a number of factors that could include the following: that there are differences in the targets of *GLK1* in *Arabidopsis* and *C. gynandra*; that a number of other transcriptional regulators are more important than *GLK1* in maintaining patterns of photosynthesis gene expression in *C. gynandra*; and that a rapid induction of *GLK1* gene expression has more impact than increasing the steady-state level of *GLK1*. This analysis is also subject to the caveat that in neither case was the amount of *GLK1* protein measured.

In all of our analyses, differences in transcript abundance between the leaves of *C. gynandra* and *C. spinosa* may reflect the operation of the C₄ and C₃ photosynthetic pathways; alternatively, they may be due to differences in metabolism and cell biology associated with the phylogenetic distance between the two species. However, in many cases, it is striking that our analysis has identified differences in the abundance of transcripts derived from genes that have been documented to be involved in processes known to alter in a C₄ leaf. Taken together, the analysis allows us to significantly extend the number of C₄-related genes controlled at the level of transcript abundance and to extend the current model for C₄-related processes in NAD-ME C₄ plants (Fig. 4). Analysis of additional pairs of C₃ and C₄ species will likely facilitate the identification of genes specifically involved in the C₄ pathway and exclude genes that are modified for other reasons.

MATERIALS AND METHODS

Plant Material and 454 Sequencing

Cleome spinosa and *Cleome gynandra* plants for transcript profiling by RNA-Seq were grown in standard potting mix in a glasshouse in August and September 2007. To obtain sequence tags for DGE analysis from *C. spinosa* and *C. gynandra*, total RNAs were isolated from fully expanded leaves sampled from 56-d-old plants of each species. mRNA was reverse transcribed to cDNA after two consecutive rounds of oligo(dT) purification and prepared for 454 sequencing as described previously (Weber et al., 2007).

Mapping and Quantification of the Sequence Reads

Evolution did not stop in the lineage to the reference genome of *Arabidopsis* (*Arabidopsis thaliana*) after the Cleomaceae branch diverged. Hence, there may be genes that were tandem duplicated or retained after the whole genome duplication event of the Brassicaceae that are absent in either of the Cleomaceae species (Bräutigam and Gowik, 2010). To avoid mapping problems such as splitting of reads or mapping errors due to differential retention of genes in either Cleomaceae or *Arabidopsis*, we created a minimal genome for mapping. The remnants of the last whole genome duplication in the lineage of the Brassicaceae (Bowers et al., 2003; Thomas et al., 2006) and the tandem duplicated genes (Haberer et al., 2004) were reduced to one representative for each based on the TAIR 9 coding sequence set. In each case, the gene with the lowest AGI code was retained for mapping. For each gene, the Supplemental Data store whether there are duplicates and which duplicates match the gene (Supplemental Tables S3 and S5). We recommend recovery of the associated duplicated genes followed by a detailed analysis with phylogenetic trees to define the true ortholog when translating the results of Cleomaceae analyses to *Arabidopsis* research.

The 454 sequence reads were mapped onto coding sequences of the minimalized TAIR 9 genome by BLAT (Kent, 2002) and BLAST (Altschul et al.,

1997) with varying parameters, and the output was parsed with in-house PERL scripts to retain only the best matching AGI codes for each sequence read and the best BLAST hit, respectively. Differentially expressed transcripts were identified using the Poisson statistics developed by Audic and Claverie (1997) followed by a Bonferroni correction to account for the accumulation of α -type errors when conducting multiple pair-wise comparisons (Audic and Claverie, 1997).

Plant Material and qPCR Analysis

Both species were grown in a growth chamber in long-day conditions (16 h of light/8 h of dark) under 350 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, at 22°C, and 65% relative humidity prior to samples being taken for qPCR. qPCR was conducted on the same samples used for RNA-Seq and also on mature leaves collected at noon grown in the growth cabinet. For qPCR, RNA was isolated using TriPure reagent (Roche Applied Science). RNA was treated with DNase I (Promega) and purified with the RNeasy Mini Kit (Qiagen). First-strand cDNA was then synthesized with SuperScriptII reverse transcriptase (Invitrogen) using 4 μg of RNA and oligo(dT) primers (Roche Applied Science). Quantitative reverse transcription-PCR was carried out with 96-well plates using a DNA Engine thermal cycler, Chromo4 real-time detector (Bio-Rad), SYBR Green JumpStart Taq Ready Mix (Sigma), and 15-fold dilution of the cDNA as a template. Initial denaturation was carried out at 94°C for 2 min, followed by 40 cycles of 94°C for 20 s, 60°C for 30 s, 72°C for 30 s, and 75°C for 5 s. Primers were designed to have melting temperatures of 60°C \pm 0.5°C and to produce amplicons of 91 to 189 bp. The specificity of the primers and lack of primer dimers in the PCR were verified using agarose gel electrophoresis and melting curve analysis. For each product, the threshold cycle CT, where the amplification reaction enters the exponential phase, was determined for three technical replicates and four independent biological replicates per species. The comparative $2^{-\Delta\Delta\text{CT}}$ method was used to quantify relative abundance of transcripts (Livak and Schmittgen, 2001). *ACTIN7* was chosen as a reference because the 454 sequencing data showed equal, intermediate levels of *ACTIN7* transcripts in both species. For the qPCR, SE values were calculated from $2^{-\Delta\Delta\text{CT}}$ values of each combination of biological replicates.

Polar Metabolite, Chlorophyll, Protein, and Enzyme Activity Analyses

For metabolite analysis, mature leaves from 56-d-old plants were collected in the middle of the light period and immediately frozen in liquid nitrogen. Three independent biological replicates were used. The tissues were ground in a mortar, and a 50-mg fresh weight aliquot was extracted using the procedure described by Lee and Fiehn (2008). Ribitol was used as an internal standard for data normalization. For GC-EI-TOF analysis, samples were processed and analyzed according to Lee and Fiehn (2008). Enzyme activities, chlorophyll, and protein content were determined according to Hausler et al. (2001).

The *Cleome* read data have been submitted to the National Center for Biotechnology Information Short Read Archive: *C. spinosa* = SRS002743.1 and *C. gynandra* = SRS002744.2.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure S1. Quantitation of marker enzyme activities in leaf extracts of *C. spinosa* and *C. gynandra*.

Supplemental Figure S2. Protein-to-fresh weight and protein-to-chlorophyll ratios in leaves of *C. gynandra* and *C. spinosa*.

Supplemental Figure S3. Changes in transcript abundance for ribosomal proteins.

Supplemental Table S1. Relative abundance of predominant metabolites detected by GC-EI-TOF in *C. gynandra* and in *C. spinosa*.

Supplemental Table S2. Number of gene loci and number of differentially expressed genes detected with BLAT and BLAST.

Supplemental Table S3. Quantitative information for all reads mapped onto the reference genome from *Arabidopsis*.

Supplemental Table S4. Comparison of mapping parameters.

Supplemental Table S5. Segmental and tandem duplicates in the *Arabidopsis* genome.

ACKNOWLEDGMENTS

We thank Tom Hardcastle for help with R.

Received May 18, 2010; accepted June 9, 2010; published June 11, 2010.

LITERATURE CITED

- Altschul SF, Madden TL, Schaffer AA, Zhang JH, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402
- Aoki N, Kanai R (1997) Reappraisal of the role of sodium in the light-dependent active transport of pyruvate into mesophyll chloroplasts of C_4 plants. *Plant Cell Physiol* 38: 1217–1225
- Aoki N, Ohnishi JI, Kanai R (1994) Proton/pyruvate cotransport into mesophyll chloroplasts of C_4 plants. *Plant Cell Physiol* 35: 801–806
- Audic S, Claverie JM (1997) The significance of digital gene expression profiles. *Genome Res* 7: 986–995
- Black CC (1973) Photosynthetic carbon fixation in relation to net CO_2 uptake. *Annu Rev Plant Physiol Plant Mol Biol* 24: 253–286
- Botha CEJ (1992) Plasmodesmatal distribution, structure and frequency in relation to assimilation in C_3 and C_4 grasses in southern Africa. *Planta* 187: 348–358
- Bowers JE, Chapman BA, Rong JK, Paterson AH (2003) Unravelling angiosperm genome evolution by phylogenetic analysis of chromosomal duplication events. *Nature* 422: 433–438
- Bräutigam A, Gowik U (2010) What can next generation sequencing do for you? Next generation sequencing as a valuable tool in plant research. *Plant Biol* 12: 831–841
- Bräutigam A, Hofmann-Benning S, Weber APM (2008a) Comparative proteomics of chloroplast envelopes from C_3 and C_4 plants reveals specific adaptations of the plastid envelope to C_4 photosynthesis and candidate proteins required for maintaining C_4 metabolite fluxes. *Plant Physiol* 148: 568–579
- Bräutigam A, Shrestha RP, Whitten D, Wilkerson CG, Carr KM, Froehlich JE, Weber APM (2008b) Comparison of the use of a species-specific database generated by pyrosequencing with databases from related species for proteome analysis of pea chloroplast envelopes. *J Biotechnol* 136: 44–53
- Bräutigam A, Weber APM (2011) Transport processes: connecting the reactions of C_4 photosynthesis. In AS Raghavendra, RF Sage, eds, *C_4 Photosynthesis and Related CO_2 Concentrating Mechanisms*. Advances in Photosynthesis and Respiration, Vol 32. Springer, Berlin, pp 199–219
- Brown NJ, Parsley K, Hibberd JM (2005) The future of C_4 research: maize, Flaveria or Cleome? *Trends Plant Sci* 10: 215–221
- Brown RH (1999) Agronomic implications of C_4 photosynthesis. In RF Sage, RK Monson, eds, *C_4 Plant Biology*. Academic Press, San Diego, pp 473–507
- Brownell PF, Crossland CJ (1972) The requirement for sodium as a micronutrient by species having the C_4 dicarboxylic photosynthetic pathway. *Plant Physiol* 49: 794–797
- Chollet R, Ogren WL (1975) Regulation of photorespiration in C_3 and C_4 species. *Bot Rev* 41: 137–179
- DeBono A, Yeats TH, Rose JKC, Bird D, Jetter R, Kunst L, Samuelsen L (2009) *Arabidopsis* LTPG is a glycosylphosphatidylinositol-anchored lipid transfer protein required for export of lipids to the plant surface. *Plant Cell* 21: 1230–1238
- De Vos M, Van Oosten VR, Van Poeckel RM, Van Pelt JA, Pozo MJ, Mueller MJ, Buchala AJ, Metraux JP, Van Loon LC, Dicke M, et al (2005) Signal signature and transcriptome changes of *Arabidopsis* during pathogen and insect attack. *Mol Plant Microbe Interact* 18: 923–937
- Dodd AN, Gardner MJ, Hotta CT, Hubbard KE, Dalchau N, Love J, Assie JM, Robertson FC, Jakobsen MK, Goncalves J, et al (2007) The *Arabidopsis* circadian clock incorporates a cADPR-based feedback loop. *Science* 318: 1789–1792
- Evert RF, Eschrich W, Heyser W (1977) Distribution and structure of plasmodesmata in mesophyll and bundle-sheath cells of *Zea mays* L. *Planta* 136: 77–89
- Fischer K, Kammerer B, Gutensohn M, Arbinger B, Weber A, Hausler RE, Flugge UI (1997) A new class of plastidic phosphate translocators: a putative link between primary and secondary metabolism by the phosphoenolpyruvate/phosphate antiporter. *Plant Cell* 9: 453–462

- Flicek P, Birney E (2009) Sense from sequence reads: methods for alignment and assembly. *Nat Methods* 6: S6–S12
- Haberer G, Hindemitt T, Meyers BC, Mayer KFX (2004) Transcriptional similarities, dissimilarities, and conservation of cis-elements in duplicated genes of *Arabidopsis*. *Plant Physiol* 136: 3009–3022
- Hatch MD (1987) C₄ photosynthesis: a unique blend of modified biochemistry, anatomy and ultrastructure. *Biochim Biophys Acta* 895: 81–106
- Hatch MD, Slack CR (1968) A new enzyme for interconversion of pyruvate and phosphopyruvate and its role in C₄ dicarboxylic acid pathway of photosynthesis. *Biochem J* 106: 141–147
- Hausler RE, Rademacher T, Li J, Lipka V, Fischer KL, Schubert S, Kreuzaler F, Hirsch HJ (2001) Single and double overexpression of C₄-cycle genes had differential effects on the pattern of endogenous enzymes, attenuation of photorespiration and on contents of UV protectants in transgenic potato and tobacco plants. *J Exp Bot* 52: 1785–1803
- Hibberd JM, Sheehy JE, Langdale JA (2008) Using C₄ photosynthesis to increase the yield of rice: rationale and feasibility. *Curr Opin Plant Biol* 11: 228–231
- Imaizumi T, Kay SA, Schroeder JI (2007) Circadian rhythms: daily watch on metabolism. *Science* 318: 1730–1731
- Jordan DB, Ogren WL (1984) The CO₂/O₂ specificity of ribulose 1,5-bisphosphate carboxylase oxygenase: dependence on ribulosebisphosphate concentration, pH and temperature. *Planta* 161: 308–313
- Keeley JE (1998) C₄ photosynthetic modifications in the evolutionary transition from land to water in aquatic grasses. *Oecologia* 116: 85–97
- Kent WJ (2002) BLAT: the BLAST-Like Alignment Tool. *Genome Res* 12: 656–664
- Kilian J, Whitehead D, Horak J, Wanke D, Wein S, Batistic O, D'Angelo C, Bornberg-Bauer E, Kudla J, Harter K (2007) The AtGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. *Plant J* 50: 347–363
- Kobayashi H, Yamada M, Taniguchi M, Kawasaki M, Sugiyama T, Miyake H (2009) Differential positioning of C₄ mesophyll and bundle sheath chloroplasts: recovery of chloroplast positioning requires the actomyosin system. *Plant Cell Physiol* 50: 129–140
- Ku MSB, Schmitt MR, Edwards GE (1979) Quantitative determination of ribulose bisphosphate carboxylase oxygenase protein in leaves of several C₃ and C₄ plants. *J Exp Bot* 114: 89–98
- Langdale JA, Kidner CA (1994) Bundle-sheath defective, a mutation that disrupts cellular differentiation in maize leaves. *Development* 120: 673–681
- Lee DY, Fiehn O (2008) High quality metabolomic data for *Chlamydomonas reinhardtii*. *Plant Methods* 4: 7
- Leegood RC (2002) C₄ photosynthesis: principles of CO₂ concentration and prospects for its introduction into C₃ plants. *J Exp Bot* 53: 581–590
- Levy A, Erlanger M, Rosenthal M, Epel BL (2007) A plasmodesmata-associated beta-1,3-glucanase in *Arabidopsis*. *Plant J* 49: 669–682
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)(-Delta Delta C) method. *Methods* 25: 402–408
- Majeran W, van Wijk KJ (2009) Cell-type-specific differentiation of chloroplasts in C₄ plants. *Trends Plant Sci* 14: 100–109
- Maple J, Fujiwara MT, Kitahata N, Lawson T, Baker NR, Yoshida S, Moller SG (2004) GIANT CHLOROPLAST 1 is essential for correct plastid division in *Arabidopsis*. *Curr Biol* 14: 776–781
- Marshall DM, Muhaidat R, Brown NJ, Liu Z, Stanley S, Griffiths H, Sage RF, Hibberd JM (2007) Cleome, a genus closely related to *Arabidopsis*, contains species spanning a developmental progression from C₃ to C₄ photosynthesis. *Plant J* 51: 886–896
- Matsuoka M, Furbank RT, Fukuyama H, Miyao M (2001) Molecular engineering of C₄ photosynthesis. *Annu Rev Plant Physiol Plant Mol Biol* 52: 297–314
- Metzker ML (2010) Applications of next-generation sequencing technologies: the next generation. *Nat Rev Genet* 11: 31–46
- Mitchell PL, Sheehy JE (2006) Supercharging rice photosynthesis to increase yield. *New Phytol* 171: 688–693
- Oaks A (1994) Efficiency of nitrogen utilization in C₃ and C₄ cereals. *Plant Physiol* 106: 407–414
- Oikawa K, Kasahara M, Kiyosue T, Kagawa T, Suetsugu N, Takahashi F, Kanegae T, Niwa Y, Kadota A, Wada M (2003) CHLOROPLAST UNUSUAL POSITIONING1 is essential for proper chloroplast positioning. *Plant Cell* 15: 2805–2815
- Osborne CP, Freckleton RP (2009) Ecological selection pressures for C₄ photosynthesis in the grasses. *Proc R Soc Lond B Biol Sci* 276: 1753–1760
- Palmieri N, Schlotterer C (2009) Mapping accuracy of short reads from massively parallel sequencing and the implications for quantitative expression profiling. *PLoS ONE* 4: 10
- Price J, Laxmi A, St Martin SK, Jang JC (2004) Global transcription profiling reveals multiple sugar signal transduction mechanisms in *Arabidopsis*. *Plant Cell* 16: 2128–2150
- Reiskind JB, Berg RH, Salvucci ME, Bowes G (1989) Immunogold localization of primary carboxylases in leaves of aquatic and a C₃-C₄ intermediate species. *Plant Sci* 61: 43–52
- Renne P, Dressen U, Hebbeker U, Hille D, Flugge UI, Westhoff P, Weber APM (2003) The *Arabidopsis* mutant *dct* is deficient in the plastidic glutamate/malate translocator DIT2. *Plant J* 35: 316–331
- Roberts AG, Oparka KJ (2003) Plasmodesmata and the control of symplastic transport. *Plant Cell Environ* 26: 103–124
- Rolland F, Baena-Gonzalez E, Sheen J (2006) Sugar sensing and signaling in plants: conserved and novel mechanisms. *Annu Rev Plant Biol* 57: 675–709
- Rossini L, Cribb L, Martin DJ, Langdale JA (2001) The maize *Golden2* gene defines a novel class of transcriptional regulators in plants. *Plant Cell* 13: 1231–1244
- Sage RF (2004) The evolution of C₄ photosynthesis. *New Phytol* 161: 341–370
- Sawers RJH, Liu P, Anufrikova K, Hwang JTG, Brutnell TP (2007) A multi-treatment experimental system to examine photosynthetic differentiation in the maize leaf. *BMC Genomics* 8: 12
- Schaffer R, Landgraf J, Accerbi M, Simon V, Larson M, Wisman E (2001) Microarray analysis of diurnal and circadian-regulated genes in *Arabidopsis*. *Plant Cell* 13: 113–123
- Schmid M, Davison TS, Henz SR, Pape UJ, Demar M, Vingron M, Scholkopf B, Weigel D, Lohmann JU (2005) A gene expression map of *Arabidopsis thaliana* development. *Nat Genet* 37: 501–506
- Sheen J (1999) C₄ gene expression. *Annu Rev Plant Physiol Plant Mol Biol* 50: 187–217
- Swarbreck D, Wilks C, Lamesch P, Berardini TZ, Garcia-Hernandez M, Foerster H, Li D, Meyer T, Muller R, Ploetz L, et al (2008) The *Arabidopsis* Information Resource (TAIR): gene structure and function annotation. *Nucleic Acids Res* 36: D1009–D1014
- Taniguchi M, Taniguchi Y, Kawasaki M, Takeda S, Kato T, Sato S, Tahata S, Miyake H, Sugiyama T (2002) Identifying and characterizing plastidic 2-oxoglutarate/malate and dicarboxylate transporters in *Arabidopsis thaliana*. *Plant Cell Physiol* 43: 706–717
- Taniguchi Y, Nagasaki J, Kawasaki M, Miyake H, Sugiyama T, Taniguchi M (2004) Differentiation of dicarboxylate transporters in mesophyll and bundle sheath chloroplasts of maize. *Plant Cell Physiol* 45: 187–200
- Thomas BC, Pedersen B, Freeling M (2006) Following tetraploidy in an *Arabidopsis* ancestor, genes were removed preferentially from one homeolog leaving clusters enriched in dose-sensitive genes. *Genome Res* 16: 934–946
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF (2005) Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of *Arabidopsis*. *Plant J* 41: 195–211
- Voznesenskaya EV, Edwards GE, Kiirats O, Artyusheva EG, Franceschi VR (2003) Development of biochemical specialization and organelle partitioning in the single-cell C₄ system in leaves of *Borszczowia aralocaspica* (Chenopodiaceae). *Am J Bot* 90: 1669–1680
- Voznesenskaya EV, Franceschi VR, Kiirats O, Artyusheva EG, Freitag H, Edwards GE (2002) Proof of C₄ photosynthesis without Kranz anatomy in *Bienertia cycloptera* (Chenopodiaceae). *Plant J* 31: 649–662
- Voznesenskaya EV, Franceschi VR, Kiirats O, Freitag H, Edwards GE (2001) Kranz anatomy is not essential for terrestrial C₄ plant photosynthesis. *Nature* 414: 543–546
- Voznesenskaya EV, Koteyeva NK, Chuong SDX, Ivanova AN, Barroca J, Craven LA, Edwards GE (2007) Physiological, anatomical and biochemical characterisation of photosynthetic types in genus *Cleome* (Cleomaceae). *Funct Plant Biol* 34: 247–267
- Wall PK, Leebens-Mack J, Chandrabali AS, Barakat A, Wolcott E, Liang HY, Landherr L, Tomsho LP, Hu Y, Carlson JE, et al (2009) Comparison

- of next generation sequencing technologies for transcriptome characterization. *BMC Genomics* **10**: 347
- Wang Z, Gerstein M, Snyder M** (2009) RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* **10**: 57–63
- Waters MT, Wang P, Korkaric M, Capper RG, Saunders NJ, Langdale JA** (2009) GLK transcription factors coordinate expression of the photosynthetic apparatus in *Arabidopsis*. *Plant Cell* **21**: 1109–1128
- Weber APM, von Caemmerer S** (2010) Plastid transport and metabolism of C₃ and C₄ plants: comparative analysis and possible biotechnological exploitation. *Curr Opin Plant Biol* **13**: 256–264
- Weber APM, Weber KL, Carr K, Wilkerson C, Ohlrogge JB** (2007) Sampling the Arabidopsis transcriptome with massively parallel pyrosequencing. *Plant Physiol* **144**: 32–42
- Wise RP, Moscou MJ, Bogdanove AJ, Whitham SA** (2007) Transcript profiling in host-pathogen interactions. *Annu Rev Phytopathol* **45**: 329–369