

# NUMERICAL APPROXIMATION OF STOCHASTIC EVOLUTION EQUATIONS: CONVERGENCE IN SCALE OF HILBERT SPACES

HAKIMA BESSAIH, ERIKA HAUSENBLAS, TSIRY RANDRIANASOLO, AND PAUL RAZAFIMANDIMBY

ABSTRACT. The present paper is devoted to the numerical approximation of an abstract stochastic nonlinear evolution equation in a separable Hilbert space  $H$ . Examples of equations which fall into our framework include the GOY and Sabra shell models and a class of nonlinear heat equations. The space-time numerical scheme is defined in terms of a Galerkin approximation in space and a semi-implicit Euler–Maruyama scheme in time. We prove the convergence in probability of our scheme by means of an estimate of the error on a localized set of arbitrary large probability. Our error estimate is shown to hold in a more regular space  $V_\beta \subset H$  with  $\beta \in [0, \frac{1}{4})$  and that the explicit rate of convergence of our scheme depends on this parameter  $\beta$ .

## 1. INTRODUCTION

Throughout this paper we fix a complete filtered probability space  $\mathfrak{U} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$  with the filtration  $\mathbb{F} = \{\mathcal{F}_t; t \geq 0\}$  satisfying the usual conditions. We also fix a separable Hilbert space  $H$  equipped with a scalar product  $(\cdot, \cdot)$  with the associated norm  $|\cdot|$  and another separable Hilbert space  $\mathcal{H}$ . In this paper, we analyze numerical approximations for an abstract stochastic evolution equation of the form

$$(1) \quad \begin{cases} d\mathbf{u} = -[A\mathbf{u} + B(\mathbf{u}, \mathbf{u})]dt + G(\mathbf{u})dW, & t \in [0, T], \\ \mathbf{u}(0) = \mathbf{u}_0, \end{cases}$$

where hereafter  $T > 0$  is a fixed number and  $A$  is a self-adjoint positive operators on  $H$ . The operators  $B$  and  $G$  are nonlinear maps satisfying several technical assumptions to be specified later and  $W = \{W(t); 0 \leq t \leq T\}$  is a  $\mathcal{H}$ -valued Wiener process.

The abstract equation (1) can describe several problems from different fields including mathematical finance, electromagnetism, and fluid dynamic. Stochastic models have been widely used to describe small fluctuations or perturbations which arise in nature. For a more exhaustive introduction to the importance of stochastic models and the analysis of stochastic partial differential equations, we refer the reader to [18, 32, 37, 40, 42].

Numerical analysis for stochastic partial differential equations (SPDEs) has known a strong interest in the past decades. Many algorithms which are based on either finite difference or finite element methods or spectral Galerkin methods (for the space discretization) and on either Euler schemes or Crank-Nicholson or Runge-Kutta schemes (for the temporal discretization) have been introduced for both the linear and nonlinear cases and their rate of convergence have been investigated widely. Here we should note that the orders of convergence that are frequently analyzed are the weak and strong orders of convergence. The literature on numerical analysis for SPDEs is now very extensive. Without being exhaustive, we only cite amongst other the recent papers [38, 16, 24, 1, 15], the excellent review paper [33] and references therein. Most of the literature deals with the stochastic heat equations with globally Lipschitz nonlinearities, but there are also several papers that treat abstract stochastic evolution equations. For example, Gyongy and Millet in [31] investigated a general evolution equation with an operator that has the strong monotone and global Lipschitz properties. They

---

*Date:* April 18, 2018.

The research of Hakima Bessaih was supported by the NSF grant DMS-1418838. The research of Erika Hausenblas was supported by the Austrian Science Fund (FWF): P25968. The research of Tsiry Randrianasolo was supported by the Austrian Science Fund (FWF): P25968 and the German Research Council as part of the Collaborative Research Center SFB 1283.

were able to implement a space-time discretization and showed a rate of convergence in mean under appropriate assumptions. Similar rate of convergence have been obtained by Bessaih and Schurz in [8] for an equation with globally Lipschitz nonlinearities. When a system of SPDEs with non-globally Lipschitz nonlinearities, such as the stochastic Navier-Stokes equations, is considered the story is completely different. Indeed, in this case the rate of convergence obtained is generally only in probability. This kind of convergence was introduced for the first time by Printems in [41] and is well suited for SPDEs with locally Lipschitz coefficients. When the stochastic perturbation is in an additive form (additive noise), then using a path wise argument one can prove a convergence in mean, we refer to Breckner in [10]. Let us mention that in this case, no rate of convergence can be deduced.

Recent literature involving nonlinear models with nonlinearities which are locally Lipschitz are [28, 17, 11, 5] and references therein. In [11] martingale solutions to the incompressible Navier-Stokes equations with Gaussian multiplicative noise are constructed from a finite element based space-time discretizations. The authors of [17] proved the convergence in probability with rates of an explicit and an implicit numerical schemes by means of a Gronwall argument. The main issue when the term  $B$  is not globally Lipschitz lies on its interplay with the stochastic forcing, which prevents a Gronwall argument in the context of expectations. This issue is for example solved in [10, 14] by the introduction of a weight, which when carefully chosen contributes in removing unwanted terms and allows to use Gronwall lemma. In [17], the authors use different approach by computing the error estimates on a sample subset  $\Omega_k \subset \Omega$  with large probability. In particular, the set  $\Omega_k$  is carefully chosen so that the random variables  $\|\nabla \mathbf{u}^\ell\|_{L^2}$  are bounded as long as the events are taken in  $\Omega_k$ , and  $\lim_{k \searrow 0} \mathbb{P}(\Omega \setminus \Omega_k) = 0$ . The result is then obtained using standard arguments based on the Gronwall lemma. Other kinds of numerical algorithms have been used in [5] for a 2D stochastic Navier-Stokes equations. There, a splitting up method has been used and a rate of convergence in probability is obtained. A blending of a splitting scheme and the method of cubature on Wiener space applied to a spectral Galerkin discretisation of degree  $N$  is used in [28] to approximate the marginal distribution of the solution of the stochastic Navier-Stokes equations on the two-dimensional torus and rates of convergence are also given. For the numerical analysis of other kind of stochastic nonlinear models that enjoy the local Lipschitz condition, without being exhaustive, we refer to [25, 26, 9, 20] and references therein. They include the stochastic Schrödinger, Burgers and KDV equations.

In the present paper, we are interested in the numerical treatment of the abstract stochastic evolution equations (1). We first give a simple and short proof of the existence and uniqueness of a mild solution and study the regularity of this solution. The result about the existence of solution is based on a fixed point argument recently developed in [12]. Then, we discretize (1) using a coupled Galerkin method and (semi-)implicit Euler scheme and show convergence in probability with rates in  $V_\beta := D(A^\beta)$ . Regarding our approach it is similar to [17] and [41], however, the results are different. Indeed, while [17] and [41] establish their rates of convergence in the space  $H$  where the solution lives, we establish our rate of convergence in  $V_\beta \subset H$  where  $\beta \in [0, \frac{1}{4})$  is arbitrary. Hence, our result does not follow from the papers [17] and [41]. In contrast to the nonlinear term of Navier-Stokes equations with periodic boundary condition treated in [17], our nonlinear term does not satisfy the property  $\langle B(\mathbf{u}, \mathbf{u}), A\mathbf{u} \rangle = 0$  which plays a crucial role in the analysis in [17]. We should also point out that our model does not fall into the general framework of the papers [31] and [8], see Remark 2.2.

Examples of semilinear equations which fall into our framework include the GOY and Sabra shell models. These toy models are used to mimic some features of turbulent flows. It seems that our work is the first one rigorously addressing the numerical approximation of such models. Our result also confirm that, in term of numerical analysis, shell models behave far better than the Navier-Stokes equations. On the theoretical point of view, we provide a new and simple proof of the existence of solutions to stochastic shell models driven by Gaussian multiplicative noise. On the physical point of view, it is also worth mentioning that shell models of turbulence are toy models which consist

of infinitely many nonlinear differential equations having a structure similar to the Fourier representation of the Navier-Stokes equations, see [27]. Moreover, they capture quite well the statistical properties of three dimensional Navier-Stokes equations, like the Kolmogorov energy spectrum and the intermittency scaling exponents for the high-order structure functions, see [27] and [29]. Due to their success in the study of turbulence, new shell models have been derived by several prominent physicists for the investigation of the turbulence in magnetohydrodynamics, see for instance [39].

Another example of system of equations which falls into our framework is a class of nonlinear heat equations described in Section 5. We do not know whether our results can cover the numerical analysis of 1D stochastic nonlinear heat equations driven by additive space-time noise. Despite this fact we believe that our paper is still interesting as we are able to treat a class of 2D stochastic nonlinear heat equations with locally Lipschitz coefficients and we are not aware of results similar to ours. In fact, most of results related to stochastic heat equations are either about 1D model, or  $d$ -dimensional,  $d \in \{1, 2, 3\}$ , models with globally Lipschitz coefficients and deal with weak convergence or convergence in weaker norm, see for instance [38, 16, 24, 1].

This paper is organized as follows: in Section 2, we introduce the necessary notations and the standing assumptions that will be used in the present work. In Section 3, we present our numerical scheme and also discuss the stability and existence of solution at each time step. The convergence of the proposed method is presented in Section 4. In Section 5 we present the stochastic shell models for turbulence and a class of stochastic nonlinear heat equations as motivating examples.

## 2. NOTATIONS, ASSUMPTIONS, PRELIMINARY RESULTS AND THE MAIN THEOREM

In this section we introduce the necessary notations and the standing assumptions that will be used in the present work. We will also introduce our numerical scheme and state our main result.

**2.1. Assumptions and notations.** Throughout this paper we fix a separable Hilbert space  $H$  with norm  $|\cdot|$  and a fixed orthonormal basis  $\{\psi_n; n \in \mathbb{N}\}$ . We assume that we are given a linear operator  $A : D(A) \subset H \rightarrow H$  which is a self-adjoint and positive operator such that the fixed orthonormal basis  $\{\psi_n; n \in \mathbb{N}\}$  satisfies

$$\{\psi_n; n \in \mathbb{N}\} \subset D(A), \quad A\psi_n = \lambda_n\psi_n,$$

for an increasing sequence of positive numbers  $\{\lambda_n; n \in \mathbb{N}\}$  with  $\lambda_n \rightarrow \infty$  as  $n \nearrow \infty$ . It is clear that  $-A$  is the infinitesimal generator of an analytic semigroup  $e^{-tA}, t \geq 0$ , on  $H$ . For any  $\alpha \in \mathbb{R}$  the domain of  $A^\alpha$  denoted by  $V_\alpha = D(A^\alpha)$  is a separable Hilbert space when equipped with the scalar product

$$(2) \quad ((\mathbf{u}, \mathbf{v}))_\alpha = \sum_{k=1}^{\infty} \lambda_k^{2\alpha} \mathbf{u}_k \mathbf{v}_k, \quad \text{for } \mathbf{u}, \mathbf{v} \in V_\alpha.$$

The norm associated to this scalar product will be denoted by  $\|\mathbf{u}\|_\alpha, \mathbf{u} \in V_\alpha$ . In what follows we set  $V := D(A^{\frac{1}{2}})$ .

Next, we consider a nonlinear map  $B(\cdot, \cdot) : V \times V \rightarrow V^*$  satisfying the following set of assumptions, where hereafter  $V^*$  denotes the dual of the Banach space  $V$ .

**(B1)** There exists a constant  $C_0 > 0$  such that for any  $\theta \in [0, \frac{1}{2})$  and  $\gamma \in (0, \frac{1}{2})$  satisfying  $\theta + \gamma \in (0, \frac{1}{2}]$ , we have

$$(3) \quad \|\mathbf{B}(\mathbf{u}, \mathbf{v}) - \mathbf{B}(\mathbf{x}, \mathbf{y})\|_{-\theta} \leq \begin{cases} C_0 \|\mathbf{u} - \mathbf{x}\|_{\frac{1}{2} - (\theta + \gamma)} (\|\mathbf{v}\|_\gamma + \|\mathbf{y}\|_\gamma) + \|\mathbf{v} - \mathbf{y}\|_\gamma (\|\mathbf{u}\|_{\frac{1}{2} - (\theta + \gamma)} + \|\mathbf{x}\|_{\frac{1}{2} - (\theta + \gamma)}) \\ \text{for any } \mathbf{u}, \mathbf{x} \in V_{\frac{1}{2} - (\theta + \gamma)} \text{ and } \mathbf{v}, \mathbf{y} \in V_\gamma, \\ C_0 (\|\mathbf{u}\|_\gamma + \|\mathbf{x}\|_\gamma) \|\mathbf{v} - \mathbf{y}\|_{\frac{1}{2} - (\theta + \gamma)} + \|\mathbf{u} - \mathbf{x}\|_\gamma (\|\mathbf{v}\|_{\frac{1}{2} - (\theta + \gamma)} + \|\mathbf{y}\|_{\frac{1}{2} - (\theta + \gamma)}) \\ \text{for any } \mathbf{v}, \mathbf{y} \in V_{\frac{1}{2} - (\theta + \gamma)} \text{ and } \mathbf{u}, \mathbf{x} \in V_\gamma. \end{cases}$$

Due to the continuous embedding  $V_{-\theta} \subset V_{-\frac{1}{2}}$ ,  $\theta \in [0, \frac{1}{2})$ , (3) holds with  $\theta$  and  $\frac{1}{2} - (\theta + \gamma)$  respectively replaced by  $\frac{1}{2}$  and  $\frac{1}{2} - \gamma$  where  $\gamma > 0$  is arbitrary.

In addition to the above, we assume that for any  $\varepsilon > 0$  there exists a constant  $C > 0$  such that

$$(4) \quad |\mathbf{B}(\mathbf{u}, \mathbf{v})| \leq C \|\mathbf{u}\| \|\mathbf{v}\|_{\frac{1}{2} + \varepsilon}, \text{ for any } \mathbf{u} \in \mathbf{H}, \mathbf{v} \in V_{\frac{1}{2} + \varepsilon}.$$

(B2) We also assume that for any  $\mathbf{u}, \mathbf{v} \in V$

$$(5) \quad \langle \mathbf{A}\mathbf{v} + \mathbf{B}(\mathbf{u}, \mathbf{v}), \mathbf{v} \rangle \geq \|\mathbf{v}\|_{\frac{1}{2}}^2.$$

(B3) We assume that for any  $\mathbf{u} \in \mathbf{H}$  we have

$$(6) \quad \mathbf{B}(0, \mathbf{u}) = \mathbf{B}(\mathbf{u}, 0) = 0.$$

Note that Assumptions (B1) and (B3) imply

(B1)' There exists a constant  $C_0 > 0$  such that for any numbers  $\theta \in [0, \frac{1}{2})$  and  $\gamma \in (0, \frac{1}{2})$  satisfying  $\theta + \gamma \in (0, \frac{1}{2}]$ , we have

$$(7) \quad \|\mathbf{B}(\mathbf{u}, \mathbf{v})\|_{-\theta} \leq C_0 \begin{cases} \|\mathbf{u}\|_{\frac{1}{2} - (\theta + \gamma)} \|\mathbf{v}\|_{\gamma} & \text{for any } \mathbf{u} \in V_{\frac{1}{2} - (\theta + \gamma)} \text{ and } \mathbf{v} \in V_{\gamma}, \\ \|\mathbf{u}\|_{\gamma} \|\mathbf{v}\|_{\frac{1}{2} - (\theta + \gamma)} & \text{for any } \mathbf{v} \in V_{\frac{1}{2} - (\theta + \gamma)}, \text{ and } \mathbf{u} \in V_{\gamma}. \end{cases}$$

If  $\theta = \frac{1}{2}$ , then (7) holds with  $\frac{1}{2} - (\theta + \gamma)$  replaced by  $\frac{1}{2} - \gamma$  where  $\gamma > 0$  is arbitrary.

Let  $\{\mathfrak{w}_j; j \in \mathbb{N}\}$  be a sequence of mutually independent and identically distributed standard Brownian motions on  $\mathfrak{U}$ . Let  $\mathcal{H}$  be separable Hilbert space and  $\mathcal{L}_1(\mathcal{H})$  be the space of all trace class operators on  $\mathcal{H}$ . Recall that if  $Q \in \mathcal{L}_1(\mathcal{H})$  is a symmetric, positive operator and  $\{\varphi_j; j \in \mathbb{N}\}$  is an orthonormal basis of  $\mathcal{H}$  consisting of eigenvectors of  $Q$ , then the series

$$W(t) = \sum_{j=1}^{\infty} \sqrt{q_j} \mathfrak{w}_j(t) \varphi_j, \quad t \in [0, T],$$

where  $\{q_j; j \in \mathbb{N}\}$  are the eigenvalues of  $Q$ , converges in  $L^2(\Omega; C([0, T]; \mathcal{H}))$  and it defines an  $\mathcal{H}$ -valued Wiener process with covariance operator  $Q$ . Furthermore, for any positive integer  $\ell > 0$  there exists a constant  $C_\ell > 0$  such that

$$(8) \quad \mathbb{E} \|W(t) - W(s)\|_{\mathcal{H}}^{2\ell} \leq C_\ell |t - s|^\ell (\text{Tr } Q)^\ell,$$

for any  $t, s \geq 0$  with  $t \neq 0$ . Before proceeding further we recall few facts about stochastic integral. Let  $\mathbf{K}$  be a separable Hilbert space,  $\mathcal{L}(\mathcal{H}, \mathbf{K})$  be the space of all bounded linear  $\mathbf{K}$ -valued operators defined on  $\mathcal{H}$ ,  $\mathcal{M}_T^2(\mathbf{K})$  be the space of all equivalence classes of  $\mathbb{F}$ -progressively measurable processes  $\Psi : \Omega \times [0, T] \rightarrow \mathbf{K}$  satisfying

$$\mathbb{E} \int_0^T \|\Psi(s)\|_{\mathbf{K}}^2 ds < \infty.$$

If  $Q \in \mathcal{L}_1(\mathcal{H})$  is a symmetric, positive and trace class operator then  $Q^{\frac{1}{2}} \in \mathcal{L}_2(\mathcal{H})$  and for any  $\Psi \in \mathcal{L}(\mathcal{H}, \mathbf{K})$  we have  $\Psi \circ Q^{\frac{1}{2}} \in \mathcal{L}_2(\mathcal{H}, \mathbf{K})$ , where  $\mathcal{L}_2(\mathcal{H}, \mathbf{K})$  (with  $\mathcal{L}_2(\mathcal{H}) := \mathcal{L}_2(\mathcal{H}, \mathcal{H})$ ) is the Hilbert space of all operators  $\Psi \in \mathcal{L}(\mathcal{H}, \mathbf{K})$  satisfying

$$\|\Psi\|_{\mathcal{L}_2(\mathcal{H}, \mathbf{K})}^2 = \sum_{j=1}^{\infty} \|\Psi \varphi_j\|_{\mathbf{K}}^2 < \infty.$$

Furthermore, from the theory of stochastic integration on infinite dimensional Hilbert space, see [21], for any  $\mathcal{L}(\mathcal{H}, \mathbf{K})$ -valued process  $\Psi$  such that  $\Psi \circ Q^{1/2} \in \mathcal{M}_T^2(\mathcal{L}_2(\mathcal{H}, \mathbf{K}))$  the process  $M$  defined by

$$M(t) = \int_0^t \Psi(s) dW(s), \quad t \in [0, T],$$

is a  $\mathbf{K}$ -valued martingale. Moreover, we have the following Itô's isometry

$$(9) \quad \mathbb{E} \left( \left\| \int_0^t \Psi(s) dW(s) \right\|_{\mathbf{K}}^2 \right) = \mathbb{E} \left( \int_0^t \|\Psi(s) Q^{\frac{1}{2}}\|_{\mathcal{L}_2(\mathcal{H}, \mathbf{K})}^2 ds \right), \quad \forall t \in [0, T],$$

and the Burkholder-Davis-Gundy inequality

$$(10) \quad \mathbb{E} \left( \sup_{0 \leq s \leq t} \left\| \int_0^s \Psi(\tau) dW(\tau) \right\|_{\mathcal{K}}^q \right) \leq C_q \mathbb{E} \left( \int_0^t \|\Psi(s) Q^{\frac{1}{2}}\|_{\mathcal{L}_2(\mathcal{H}, \mathcal{K})}^2 ds \right)^{\frac{q}{2}}, \forall t \in [0, T], \forall q \in (1, \infty).$$

Now, we impose the following set of conditions on the nonlinear term  $G(\cdot)$  and the Wiener process  $W$ .

- (N) Let  $\mathcal{H}$  be a separable Hilbert space. We assume that the driving noise  $W$  is a  $\mathcal{H}$ -valued Wiener process with a positive and symmetric covariance operator  $Q \in \mathcal{L}_1(\mathcal{H})$ .
- (G) We assume that the nonlinear function  $G : \mathbb{H} \rightarrow \mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})$  is measurable and that there exists a constant  $C_1 > 0$  such that for any  $\mathbf{u} \in \mathbb{H}, \mathbf{v} \in \mathbb{H}$  we have

$$\|G(\mathbf{u}) - G(\mathbf{v})\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})} \leq C_1 |\mathbf{u} - \mathbf{v}|.$$

**Remark 2.1.**

- (a) Note that the above assumption implies that  $G : \mathbb{H} \rightarrow \mathcal{L}(\mathcal{H}, \mathbb{H})$  is globally Lipschitz and of at most linear growth, i.e, there exists a constant  $C_2 > 0$  such that

$$\begin{aligned} \|G(\mathbf{u}) - G(\mathbf{v})\|_{\mathcal{L}(\mathcal{H}, \mathbb{H})} &\leq C_2 |\mathbf{u} - \mathbf{v}|, \\ |G(\mathbf{u})| &\leq C_2 (1 + |\mathbf{u}|), \end{aligned}$$

for any  $\mathbf{u}, \mathbf{v} \in \mathbb{H}$ .

- (b) There also exists a number  $C_3 > 0$  such that

$$\begin{aligned} \|G(\mathbf{u}) - G(\mathbf{v})\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})} &\leq C_3 \|\mathbf{u} - \mathbf{v}\|_{\frac{1}{4}}, \\ \|G(\mathbf{u})\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})} &\leq C_3 (1 + \|\mathbf{u}\|_{\frac{1}{4}}), \end{aligned}$$

for any  $\mathbf{u}, \mathbf{v} \in V_{\frac{1}{4}}$ .

- (c) Owing to item (a) of the present remark, if  $\mathbf{u} \in \mathcal{M}_T^2(\mathbb{H})$ , then  $G(\mathbf{u}) \circ Q^{\frac{1}{2}} \in \mathcal{M}_T^2(\mathcal{L}_2(\mathcal{H}, \mathbb{H}))$  and the stochastic integral  $\int_0^t G(\mathbf{u}(s)) dW(s)$  is a well defined  $\mathbb{H}$ -valued martingale.

To close the current subsection we formulate the following remark.

**Remark 2.2.** Our assumptions on our problem do not imply the assumptions in either [31] or [8]. To justify this claim assume that the coefficient of the noise  $G$  of our paper and those of [31] and [8] are both zero. Let us now set

$$A(t, u) = -Au - B(u, u),$$

which basically corresponds to the drift in both [31] and [8]. For the sake of simplicity we take  $\theta = 0$  and  $\gamma = \frac{1}{4}$  in our assumption (B1). The spaces  $H$  and  $V$  in [31] and [8] are respectively  $V_0$  and  $V_{\frac{1}{2}}$  in our framework. The map  $A(t, u)$  defined above satisfies

$$\langle A(t, u) - A(t, v), u - v \rangle \leq -|u - v|^2 + C_0 |u - v| \|u - v\|_{\frac{1}{4}} \left( \|u\|_{\frac{1}{4}} + \|v\|_{\frac{1}{4}} \right).$$

This implies that our assumptions does not imply either [31, Assumptions 2.1(i) and (2.2)(1)] or [8, Assumption (H2)].

**2.2. Preliminary results.** In this subsection we recall and derive some results that will be used in the remaining part of the paper. To this end, we first define the notion of solution of (1).

**Definition 2.3.** An  $\mathbb{F}$ -adapted process  $\mathbf{u}$  is called a weak solution of (1) (in the sense of PDEs) if the following conditions are satisfied

- (i)  $\mathbf{u} \in L^2(0, T; V) \cap C([0, T]; \mathbb{H})$   $\mathbb{P}$ -a.s.,
- (ii) for every  $t \in [0, T]$  we have  $\mathbb{P}$ -a.s.

$$(11) \quad (\mathbf{u}(t), \phi) = (\mathbf{u}_0, \phi) - \int_0^t (\langle A\mathbf{u}(s) + \mathbf{B}(\mathbf{u}(s), \mathbf{u}(s)), \phi \rangle) ds + \int_0^t \langle \phi, G(\mathbf{u}(s)) dW(s) \rangle,$$

for any  $\phi \in V$ .

**Definition 2.4.** An  $\mathbb{F}$ -adapted process  $\mathbf{u} \in C([0, T]; \mathbb{H})$   $\mathbb{P}$ -a.s. is called a mild solution to (1) if for every  $t \in [0, T]$ ,

$$(12) \quad \mathbf{u}(t) = e^{-tA}\mathbf{u}_0 + \int_0^t e^{-(t-r)A}\mathbf{B}(\mathbf{u}(r), \mathbf{u}(r))dr + \int_0^t e^{-(t-r)A}\mathbf{G}(\mathbf{u}(r))d\mathbf{W}(r), \mathbb{P}\text{-a.s.}$$

**Remark 2.5.** Observe that if  $\mathbf{u} \in L^2(0, T; \mathbb{V}) \cap C([0, T], \mathbb{H})$  is a mild solution to (1), then for any  $t > s \geq 0$ ,

$$\mathbf{u}(t) = e^{-(t-s)A}\mathbf{u}(s) + \int_s^t e^{-(t-r)A}\mathbf{B}(\mathbf{u}(r), \mathbf{u}(r))dr + \int_s^t e^{-(t-r)A}\mathbf{G}(\mathbf{u}(r))d\mathbf{W}(r), \mathbb{P}\text{-a.s.}$$

In fact, we have

$$\begin{aligned} \mathbf{u}(t) &= e^{-(t-s)A} \left( e^{-sA}\mathbf{u}_0 + \int_0^s e^{-(s-r)A}\mathbf{B}(\mathbf{u}(r), \mathbf{u}(r))dr + \int_0^s e^{-(s-r)A}\mathbf{G}(\mathbf{u}(r))d\mathbf{W}(r) \right) \\ &\quad + \int_s^t e^{-(t-r)A}\mathbf{B}(\mathbf{u}(r), \mathbf{u}(r))dr + \int_s^t e^{-(t-r)A}\mathbf{G}(\mathbf{u}(r))d\mathbf{W}(r) \\ &= e^{-(t-s)A}\mathbf{u}(s) + \int_s^t e^{-(t-r)A}\mathbf{B}(\mathbf{u}(r), \mathbf{u}(r))dr + \int_s^t e^{-(t-r)A}\mathbf{G}(\mathbf{u}(r))d\mathbf{W}(r), \mathbb{P}\text{-a.s.} \end{aligned}$$

This remark is used later to prove a very important lemma for our analysis, see Lemma 4.1.

Next, we state and give a short proof of the following results.

**Proposition 2.6.** If the assumptions (B1) to (B3) hold and (G) is satisfied with  $\mathbb{V}_{\frac{1}{4}}$  replaced by  $\mathbb{H}$  and  $\mathbf{u}_0 \in L^2(\Omega, \mathbb{H})$ , then the problem (1) has a unique global mild, which is also a weak, solution  $\mathbf{u}$ . Moreover, if  $\mathbf{u}_0 \in L^{2p}(\Omega, \mathbb{H})$  for any real number  $p \in [2, 8]$ , then there exists a constant  $\mathcal{C} > 0$  such that

$$(13) \quad \mathbb{E} \sup_{t \in [0, T]} |\mathbf{u}(t)|^{2p} + \mathbb{E} \int_0^T |\mathbf{u}(s)|^{2p-2} |A^{\frac{1}{2}}\mathbf{u}(s)|^2 ds \leq \mathcal{C}(1 + \mathbb{E}|\mathbf{u}_0|^{2p}),$$

and

$$(14) \quad \mathbb{E} \left( \int_0^T |A^{\frac{1}{2}}\mathbf{u}(s)|^2 ds \right)^p \leq \mathcal{C}(1 + \mathbb{E}|\mathbf{u}_0|^{2p}).$$

If, in addition, Assumption (G) is satisfied and  $\mathbf{u}_0 \in L^p(\Omega, \mathbb{V}_{\frac{1}{4}})$  with  $p \in [2, 8]$ , then there exists a constant  $\mathcal{C} > 0$  such that

$$(15) \quad \mathbb{E} \sup_{t \in [0, T]} \|\mathbf{u}(t)\|_{\frac{1}{4}}^p + \mathbb{E} \left( \int_0^T \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 ds \right)^p \leq \mathcal{C}(1 + \mathbb{E}\|\mathbf{u}_0\|_{\frac{1}{4}}^p + (\mathbb{E}|\mathbf{u}_0|^{2p})^2).$$

*Proof.* Let us first prove the existence of a local mild solution. For this purpose, we study the properties of  $\mathbf{B}$  in order to apply a contraction principle as in [12, Theorem 3.15]. Let  $\mathbf{B}(\cdot)$  be the mapping defined by  $\mathbf{B}(\mathbf{x}) = \mathbf{B}(\mathbf{x}, \mathbf{x})$  for any  $\mathbf{x} \in \mathbb{V}_\beta$ . Let  $\beta \in (0, \frac{1}{2})$ . Using Assumptions (B1) with  $\theta = \frac{1}{2} - \beta$ ,  $\gamma = \beta$ , we derive that

$$(16) \quad \|\mathbf{B}(\mathbf{x}) - \mathbf{B}(\mathbf{y})\|_{\beta - \frac{1}{2}} \leq C_0 \|\mathbf{x} - \mathbf{y}\| (\|\mathbf{x}\|_\beta + \|\mathbf{y}\|_\beta) + C \|\mathbf{x} - \mathbf{y}\|_\beta (\|\mathbf{x}\| + \|\mathbf{y}\|),$$

for any  $\mathbf{x}, \mathbf{y} \in \mathbb{V}_\beta$ . Since, by [43, Theorem 1.18.10, pp 141],  $\mathbb{V}_\beta$  coincides with the complex interpolation  $[\mathbb{H}, D(A^{\frac{1}{2}})]_{2\beta}$ , we infer from the interpolation inequality [43, Theorem 1.9.3, pp 59] and (16) that

$$(17) \quad \|\mathbf{B}(\mathbf{x}) - \mathbf{B}(\mathbf{y})\|_{\beta - \frac{1}{2}} \leq C_0 \|\mathbf{x} - \mathbf{y}\| (|\mathbf{x}|^{1-2\beta} \|\mathbf{x}\|_{\frac{1}{2}}^{2\beta} + |\mathbf{y}|^{1-2\beta} \|\mathbf{y}\|_{\frac{1}{2}}^{2\beta}) + C \|\mathbf{x} - \mathbf{y}\|_{\frac{1}{2}}^{2\beta} |\mathbf{x} - \mathbf{y}|^{1-2\beta} (\|\mathbf{x}\| + \|\mathbf{y}\|),$$

for any  $\mathbf{x}, \mathbf{y} \in \mathbb{V}$ . Now, we denote by  $X_T$  the Banach space  $C([0, T]; \mathbb{H}) \cap L^2(0, T; \mathbb{V})$  endowed with the norm

$$\|\mathbf{x}\|_{X_T} = \sup_{t \in [0, T]} |\mathbf{x}(t)| + \left( \int_0^T \|\mathbf{x}(t)\|_{\frac{1}{2}}^2 dt \right)^{\frac{1}{2}}.$$

We recall the following classical result, see [22, Theorem 3, pp 520].

$$(18) \quad \text{The linear map } \Lambda : L^2(0, T; \mathbb{V}^*) \ni f \mapsto \mathbf{x}(\cdot) = \int_0^\cdot e^{-(\cdot-r)A} f(r) dr \in X_T \text{ is continuous.}$$

Thus, thanks to (17), (18) and Assumption (G) we can apply [12, Theorem 3.15] to infer the existence of a unique local mild solution  $\mathbf{u}$  with lifespan  $\tau$  of (1) (we refer to [12, Definition 3.1] for the definition of local solution). Let  $\{\tau_j; j \in \mathbb{N}\}$  be an increasing sequence of stopping times converging almost surely to the lifespan  $\tau$ . Using the equivalence lemma in [21, Proposition 6.5] we can easily prove that the local mild solution is also a local weak solution satisfying (11) with  $t$  replaced by  $t \wedge \tau_j$ ,  $j \in \mathbb{N}$ . Now, we can prove by arguing as in [13, Appendix A] or [14, Proof of Theorem 4.4] that the local solution  $\mathbf{u}$  satisfies (13) uniformly w.r.t.  $j \in \mathbb{N}$ . With this observation along with an argument similar to [12, Proof of Theorem 2.10] we conclude that (1) admits a global solution (*i.e.*,  $\tau = T$  a.s.)  $\mathbf{u}$  satisfying (13) and  $\mathbf{u} \in X_T$  almost-surely.

As mentioned earlier the proof follows a similar argument as in [13, Appendix A], but for the sake of completeness we sketch the proof of (13). We apply Itô's formula first to  $|\cdot|$  and the process  $\mathbf{u}(\cdot \wedge \tau_j)$  and then to the map  $x \rightarrow x^p$   $p \geq 2$  and the process  $|\mathbf{u}(\cdot \wedge \tau_j)|^2$ . Then, using the assumption (B2) and (G) we infer that there exists a constant  $\mathcal{C} > 0$  such that for any  $j \in \mathbb{N}$

$$\begin{aligned} & \sup_{t \in [0, T]} |\mathbf{u}(t \wedge \tau_j)|^{2p} + \int_0^T |\mathbf{u}(s)|^{2p-2} |A^{\frac{1}{2}} \mathbf{u}(s)|^2 ds \leq \mathcal{C} \mathbb{E} |\mathbf{u}_0|^{2p} \\ & + \mathcal{C} \int_0^T |\mathbf{u}(s \wedge \tau_j)|^{2p-2} |(1 + |\mathbf{u}(s \wedge \tau_j)|^2)|^2 ds \\ & + 2p \sup_{t \in [0, T]} \int_0^{t \wedge \tau_j} |\mathbf{u}(s)|^{2p-2} \langle \mathbf{u}(s), G(\mathbf{u}(s)) dW(s) \rangle. \end{aligned}$$

Using the Burkholder-Holder-Davis inequality we deduce that

$$\begin{aligned} \mathbb{E} \sup_{t \in [0, T]} \int_0^{t \wedge \tau_j} |\mathbf{u}(s)|^{2p-2} \langle G(\mathbf{u}(s)), \mathbf{u}(s) \rangle dW(s) & \leq \mathbb{E} \left( \int_0^T (|\mathbf{u}(s \wedge \tau_j)|^{4p} ds) \right)^{1/2} \\ & + \mathbb{E} \left( \int_0^T (|\mathbf{u}(s \wedge \tau_j)|^{4p-2} ds) \right)^{1/2}. \end{aligned}$$

Using Young's inequality, we infer that for any  $\epsilon \in (0, \frac{1}{2})$  there exists a constant  $C(\epsilon) > 0$  such that

$$\mathbb{E} \left( \int_0^T (|\mathbf{u}(s \wedge \tau_j)|^{4p} ds) \right)^{1/2} \leq \epsilon \mathbb{E} \sup_{t \in [0, T]} |\mathbf{u}(t \wedge \tau_j)|^{2p} + C(\epsilon) \mathbb{E} \int_0^T \sup_{s \in [0, t]} |\mathbf{u}(s \wedge \tau_j)|^{2p} dt.$$

For the second integral, we need to use Hölder's inequality and then Young's inequality and the previous calculations

$$\mathbb{E} \left( \int_0^T (|\mathbf{u}(s \wedge \tau_j)|^{4p-2} ds) \right)^{1/2} \leq \epsilon \mathbb{E} \sup_{t \in [0, T]} |\mathbf{u}(t \wedge \tau_j)|^{2p} + C(\epsilon) \mathbb{E} \int_0^T \sup_{s \in [0, t]} |\mathbf{u}(s \wedge \tau_j)|^{2p} dt + T^{\frac{1}{2p}}.$$

Now collecting all the estimates we get that

$$\begin{aligned} (1 - 2\epsilon) \mathbb{E} \sup_{t \in [0, T]} |\mathbf{u}(t \wedge \tau_j)|^{2p} + \int_0^T \mathbb{E} |\mathbf{u}(s)|^{2p-2} |A^{\frac{1}{2}} \mathbf{u}(s)|^2 ds & \leq \mathcal{C} (1 + \mathbb{E} |\mathbf{u}_0|^{2p}) \\ & + \mathcal{C} \mathbb{E} \int_0^T \sup_{s \in [0, t]} |\mathbf{u}(s \wedge \tau_j)|^{2p} dt. \end{aligned}$$

Now, choosing  $\epsilon = \frac{1}{4}$ , applying Gronwall's lemma and passing to the limit as  $j \rightarrow \infty$  complete the proof of (13). The estimate (14) easily follows from (13), so we omit its proof.

We shall now prove the inequality (15). To start with we will apply Itô's formula to  $\varphi(\mathbf{u}) = \|\mathbf{u}\|_{\frac{1}{4}}^2$ . Note that thanks to the estimates (13) and (14), Assumptions (B1) and (G) we readily check that there exists a constant  $C > 0$  such that

$$\mathbb{E} \int_0^T \left[ \|A\mathbf{u} + B(\mathbf{u}, \mathbf{u})\|_{-\frac{1}{2}}^2 + \|G(\mathbf{u})\|_{\mathcal{L}(\mathfrak{H}, V_{\frac{1}{4}})}^2 \right] (t) dt \leq C.$$

Hence the general Itô's formula in [36, Section 3] is applicable to (1) and the functional  $\varphi(\mathbf{u})(t) = \|\mathbf{u}(t)\|_{\frac{1}{4}}^2$ . Thus, an application of Itô's formula to the functional  $\varphi(\mathbf{u})(t \wedge \tau_j) = \|\mathbf{u}(t \wedge \tau_j)\|_{\frac{1}{4}}^2$  gives

$$\varphi(\mathbf{u}(t \wedge \tau_j)) = \varphi(\mathbf{u}(0)) + \int_0^{t \wedge \tau_j} \varphi'(\mathbf{u}(s)) d\mathbf{u}(s) + \frac{1}{2} \int_0^{t \wedge \tau_j} \text{Tr}(\varphi''(\mathbf{u}(s))G(\mathbf{u}(s))Q(G\mathbf{u}(s))^*) ds,$$

which along with the inequality  $\frac{1}{2}\|\varphi''(\mathbf{u})\| \leq 1$ , where the norm is understood as the norm of a bilinear map, implies

$$(19) \quad \begin{aligned} & \|\mathbf{u}(t \wedge \tau_j)\|_{\frac{1}{4}}^2 + 2 \int_0^{t \wedge \tau_j} \left( \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 + 2\langle A^{\frac{1}{2}}\mathbf{u}(s), B(\mathbf{u}(s), \mathbf{u}(s)) \rangle \right) ds \\ & \leq \|\mathbf{u}_0\|_{\frac{1}{4}}^2 + 2 \int_0^{t \wedge \tau_j} \langle A^{\frac{1}{2}}\mathbf{u}(s), G(\mathbf{u}(s))dW(s) \rangle + C \text{Tr} Q \int_0^{t \wedge \tau_j} \|G(\mathbf{u}(s))\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 ds. \end{aligned}$$

Since the embedding  $V_{\frac{1}{2}+\alpha} \subset V_{2\alpha}$  is continuous for any  $\alpha \in [0, \frac{1}{2}]$ , we can use Assumptions (B1)' and the Cauchy inequality to infer that

$$\begin{aligned} \left| \int_0^{t \wedge \tau_j} \langle A^{\frac{1}{2}}\mathbf{u}(s), B(\mathbf{u}(s), \mathbf{u}(s)) \rangle ds \right| & \leq C \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{2}} \|B(\mathbf{u}(s), \mathbf{u}(s))\| ds, \\ & \leq \frac{1}{2} \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 ds + C \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{2}-\gamma}^2 \|\mathbf{u}(s)\|_{\gamma}^2 ds, \end{aligned}$$

for some  $\gamma \in (0, \frac{1}{2})$ . From an application of a complex interpolation inequality, see [43, Theorem 1.9.3, pp 59], we infer that

$$\left| \int_0^T \langle A^{\frac{1}{2}}\mathbf{u}(s), B(\mathbf{u}(s), \mathbf{u}(s)) \rangle ds \right| \leq \frac{1}{2} \int_0^T \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 ds + \int_0^T |\mathbf{u}(s)|^2 \|\mathbf{u}(s)\|_{\frac{1}{2}}^2 ds.$$

Plugging the latter inequality into (19), using the assumption on  $G$  we obtain

$$(20) \quad \begin{aligned} \|\mathbf{u}(t \wedge \tau_j)\|_{\frac{1}{4}}^2 + \frac{3}{2} \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 ds & \leq \|\mathbf{u}(0)\|_{\frac{1}{4}}^2 + C \sup_{s \in [0, T]} |\mathbf{u}(s)|^2 \int_0^T \|\mathbf{u}(s)\|_{\frac{1}{2}}^2 ds \\ & + CT + C \int_0^T \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds + 2 \left| \int_0^{t \wedge \tau_j} \langle A^{\frac{1}{4}}\mathbf{u}(s), A^{\frac{1}{4}}G(\mathbf{u}(s))dW(s) \rangle \right|. \end{aligned}$$

Taking the supremum over  $t \in [0, T]$ , then raising both sides of the resulting inequality to the power  $p/2$ , taking the mathematical expectation, and finally using the Burkholder-Davis-Gundy inequality yield

$$(21) \quad \begin{aligned} \mathbb{E} \sup_{s \in [0, t]} \|\mathbf{u}(s \wedge \tau_j)\|_{\frac{1}{4}}^p + 2\mathbb{E} \left( \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 ds \right)^{p/2} & - \left( C\mathbb{E}\|\mathbf{u}(0)\|_{\frac{1}{4}}^p + CT + C\mathbb{E} \left[ \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds \right]^{\frac{p}{2}} \right) \\ & \leq C \left( \mathbb{E} \sup_{s \in [0, T]} |\mathbf{u}(s)|^{2p} \right)^{\frac{1}{2}} \left[ \mathbb{E} \left( \int_0^T \|\mathbf{u}(s)\|_{\frac{1}{2}}^2 ds \right)^p \right]^{\frac{1}{2}} \\ & + 2C\mathbb{E} \left( \int_0^{t \wedge \tau_j} |A^{\frac{1}{4}}\mathbf{u}(s)|^2 \|G(\mathbf{u}(s))\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 ds \right)^{\frac{p}{4}}. \end{aligned}$$

Here we have used the fact that for any integer  $\ell$  and  $n$  we can find a constant  $C_{\ell, n}$  such that

$$(22) \quad \sum_{i=1}^n a_i^\ell \leq \left( \sum_{i=1}^n a_i \right)^\ell \leq C_{\ell, n} \sum_{i=1}^n a_i^\ell$$

for a sequence of non-negative numbers  $\{a_i; i = 1, 2, \dots, n\}$ .



Using the assumptions on  $G$  and Young's inequality we infer that there exists a constant  $C > 0$  such that for any  $j \in \mathbb{N}$

$$\mathbb{E} \left( \int_0^{t \wedge \tau_j} |A^{\frac{1}{4}} \mathbf{u}(s)|^2 \|G(\mathbf{u}(s))\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 ds \right)^{\frac{p}{4}} \leq \frac{1}{4} \mathbb{E} \sup_{s \in [0, t]} \|\mathbf{u}(s \wedge \tau_j)\|_{\frac{1}{4}}^p + C \mathbb{E} \left[ \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds \right]^{\frac{p}{2}} + CT,$$

which along with (21), (13) and (14) implies

$$\begin{aligned} \mathbb{E} \sup_{s \in [0, t]} \|\mathbf{u}(s \wedge \tau_j)\|_{\frac{1}{4}}^p + 2 \mathbb{E} \left( \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds \right)^{\frac{p}{2}} &\leq \mathbb{E} \|\mathbf{u}(0)\|_{\frac{1}{4}}^p + C^2 (1 + \mathbb{E} |\mathbf{u}_0|^{2p})^2 \\ &\quad + CT + \mathbb{E} \left[ \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds \right]^{\frac{p}{2}}. \end{aligned}$$

Now, we infer from the interpolation inequality [43, Theorem 1.9.3, pp 59], (13) and (14) that there exists a constant  $C > 0$  such that for any  $j \in \mathbb{N}$

$$\begin{aligned} \mathbb{E} \left[ \int_0^{t \wedge \tau_j} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 ds \right]^{\frac{p}{2}} &\leq T^{\frac{p}{2}} \mathbb{E} \left( \sup_{s \in [0, T]} |\mathbf{u}(s)|^{\frac{p}{2}} \left[ \int_0^T \|\mathbf{u}(s)\|^2 ds \right]^{\frac{p}{4}} \right) \\ &\leq CT. \end{aligned}$$

Hence,

$$\mathbb{E} \sup_{s \in [0, t]} \|\mathbf{u}(s \wedge \tau_j)\|_{\frac{1}{4}}^p \leq C_T (1 + \mathbb{E} \|\mathbf{u}(0)\|_{\frac{1}{4}}^p + (\mathbb{E} |\mathbf{u}_0|^{2p})^2),$$

from which along with a passage to the limit we readily complete the proof of the proposition.  $\square$

**2.3. The numerical scheme and the main result.** Let  $N$  be a positive integer,  $H_N \subset H$  the linear space spanned by  $\{\psi_n; n = 1, \dots, N\}$ , and  $\pi_N : H \rightarrow H_N$  the orthogonal projection of  $H$  onto the finite dimensional subspace  $H_N$ . The projection of  $\mathbf{u}$  by  $\pi_N$  is denoted by

$$(23) \quad \mathbf{u}^N := \pi_N \mathbf{u} = \sum_{n=1}^N (\psi_n, \mathbf{u}) \psi_n,$$

for  $\mathbf{u} \in H$ . The Galerkin approximation of the SPDEs (1) reads

$$(24) \quad d\mathbf{u}^N = [\pi_N A \mathbf{u}^N + \pi_N B(\mathbf{u}^N, \mathbf{u}^N)] dt + \pi_N G(\mathbf{u}^N) dW(t), \quad \mathbf{u}^N(0) = \pi_N \mathbf{u}_0.$$

Due to the assumptions (B1)-(B3) and (G), we can use Proposition 2.6 to prove that (24) has a global weak solution.

To derive an approximation of the exact solution  $\mathbf{u}$  of (1) we construct an approximation  $U^j$  of the Galerkin solution  $\mathbf{u}^N$ . To this end, let  $M$  be a positive integer and  $I_M = ([t_m, t_{m+1}))_{m=0}^M$  an equidistant grid of mesh-size  $k = t_{m+1} - t_m$  covering  $[0, T]$ . Now, for any  $j \in \{0, \dots, M-1\}$  we look for a sequence of  $\mathbb{F}$ -adapted random variables  $U^j \in H_N, j = 0, 1, \dots, M$  such that for any  $w \in V$

$$(25) \quad \begin{cases} U^0 = \pi_N \mathbf{u}_0, \\ \langle U^{j+1} - U^j + k[\pi_N A U^{j+1} + \pi_N B(U^j, U^{j+1})], w \rangle = \langle w, \pi_N G(U^j) \Delta_{j+1} W \rangle, \end{cases}$$

where  $\Delta_{j+1} W := W(t_{j+1}) - W(t_j), j \in \{0, \dots, M-1\}$ , is an independent and identically distributed random variables. We will justify in the following proposition that for a given  $U_0 = \pi_N \mathbf{u}_0$  the numerical scheme (25) admits at least one solution  $U^j \in H_N, j \in \{1, \dots, M\}$  and that (25) is stable in  $H$  and  $D(A^{\frac{1}{4}})$ .

**Proposition 2.7.** *Let the assumptions (B1)-(B3) and (G) hold. Let  $N$  and  $M$  be two fixed positive integers and  $\mathbf{u}_0 \in L^{2p}(\Omega; H)$  for any integer  $p \in [2, 4]$ . Then, for any  $j \in \{1, \dots, M\}$  there exists at least a  $\mathcal{F}_{t_j}$ -measurable*

random variable  $U^j \in H_N$  satisfying (25). Moreover, there exists a constant  $C > 0$  (depending only on  $T$  and  $\text{Tr } Q$ ) such that

$$(26) \quad \mathbb{E} \max_{0 \leq m \leq M} |U^m|^2 + \sum_{j=0}^{M-1} |U^{j+1} - U^j|^2 + 2k \mathbb{E} \sum_{j=1}^M \|U^j\|_{\frac{1}{2}}^2 \leq C(\mathbb{E}|\mathbf{u}_0|^2 + 1),$$

$$(27) \quad \mathbb{E} \left[ \max_{1 \leq m \leq M} |U^m|^{2p} + k \sum_{j=1}^M |U^j|^{2p-1} \|U^j\|_{\frac{1}{2}}^2 \right] \leq C(1 + \mathbb{E}|\mathbf{u}_0|^{2p-1}),$$

$$(28) \quad \mathbb{E} \left[ k \sum_{j=1}^M \|U^j\|_{\frac{1}{2}}^2 \right]^{2p-1} \leq C(1 + \mathbb{E}|\mathbf{u}_0|^{2p}).$$

Furthermore, if  $\mathbf{u}_0 \in L^8(\Omega, D(A^{\frac{1}{4}}))$ , then there exists a constant  $C > 0$  such that

$$(29) \quad \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^2 + \mathbb{E} \sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + k \mathbb{E} \sum_{j=1}^M \|U^j\|_{\frac{3}{4}}^2 \leq C,$$

$$(30) \quad \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^4 + \mathbb{E} \left( \sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 \right)^2 + k^2 \mathbb{E} \left( \sum_{j=1}^M \|U^j\|_{\frac{3}{4}}^2 \right)^2 \leq C$$

*Proof.* The detailed proofs of the existence, measurability and the estimates (29) and (30) will be given in Section 3. Thanks to the assumption (B2), the proof of the inequalities (26)-(28) is very similar to the proof of [17], so we omit it.  $\square$

We should note that the estimates (29) and (30) hold even if  $\mathbf{u}_0 \in L^4(\Omega, D(A^{\frac{1}{4}}))$ , but for the sake of consistency we take  $\mathbf{u}_0 \in L^8(\Omega, D(A^{\frac{1}{4}}))$ .

Now, we proceed to the statement of the main result of this paper.

**Theorem 2.8.** *Let the assumptions (B1)-(B3) and (G) hold and assume that  $\mathbf{u}_0 \in L^{16}(\Omega; H) \cap L^8(\Omega; V_{\frac{1}{4}})$ . Then for any  $\beta \in [0, \frac{1}{4})$ , there exists a constant  $k_0 > 0$  such that for any small number  $\varepsilon > 0$  we have*

$$(31) \quad \max_{1 \leq j \leq M} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \|\mathbf{u}(t_j) - U^j\|_{\beta}^2 \right) + 2k \mathbb{E} \left( \mathbf{1}_{\Omega_k} \sum_{j=1}^M \|\mathbf{u}(t_j) - U^j\|_{\frac{1}{2}+\beta}^2 \right) < k_0 k^{-2\varepsilon} [k^{2(\frac{1}{4}-\beta)} + \Lambda_N^{-2(\frac{1}{4}-\beta)}],$$

where the set  $\Omega_k$  is defined by

$$\Omega_k = \left\{ \omega \in \Omega : \sup_{t \in [0, T]} \|\mathbf{u}(t, \omega)\|_{\frac{1}{4}}^2 < \log k^{-\varepsilon}, \max_{0 \leq j \leq M} \|U^j(\omega)\|_{\frac{1}{4}}^2 < \log k^{-\varepsilon} \right\}.$$

*Proof.* The proof of this theorem will be given in Section 4.  $\square$

**Remark 2.9.** *Note that owing to (15) and (30) and the Markov inequality it is not difficult to prove that the set  $\Omega_k$  satisfies*

$$\lim_{k \searrow 0} \mathbb{P}[\Omega \setminus \Omega_k] = 0.$$

**Corollary 2.10.** *If all the assumptions of Theorem 2.8 are satisfied, then the solution  $\{U^j; j = 1, 2, \dots, M\}$  of the numerical scheme (25) converges in probability in the Hilbert space  $V_{\beta}$ ,  $\beta \in [0, \frac{1}{4})$ . More precisely, for any small number  $\varepsilon > 0$ , any  $\theta_0 \in (0, \frac{1}{4} - \beta - \varepsilon)$  and  $\theta_1 \in (0, \frac{1}{4} - \beta)$  we have*

$$(32) \quad \lim_{\Theta \nearrow \infty} \lim_{k \searrow 0} \lim_{N \nearrow \infty} \max_{1 \leq j \leq M} \mathbb{P} \left( \|\mathbf{u}(t_j) - U^j\|_{\beta} + k^{\frac{1}{2}} \left( \sum_{j=1}^M \|\mathbf{u}(t_j) - U^j\|_{\frac{1}{2}+\beta}^2 \right)^{\frac{1}{2}} \geq \Theta [k^{\theta_0} + \Lambda_N^{-\theta_1}] \right) = 0.$$

*Proof.* To shorten notation let us set  $\mathbf{e}^j := \mathbf{u}(t_j) - U^j$  and

$$\Omega_{k,N}^{\Theta} = \left\{ \omega \in \Omega; \|\mathbf{e}^j\|_{\beta}^2 + k \sum_{j=1}^M \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \geq \Theta [k^{\theta_0} + \Lambda_N^{-\theta_1}] \right\},$$

for any positive numbers  $M$  and  $k$ . Let  $\Omega_k$  be as in the statement of Theorem 2.8. Owing to (31), (15), (30) and the Chebychev-Markov inequality, we can find a constant  $\tilde{C}_5 > 0$  such that

$$\begin{aligned} \mathbb{P}\left(\Omega_{k,N}^\ominus\right) &= \mathbb{P}\left(\Omega_{k,N}^\ominus \cap \Omega_k\right) + \mathbb{P}\left(\Omega_{k,N}^\ominus \cap \Omega_k^c\right) \\ &\leq \mathbb{P}\left(\Omega_{k,N}^\ominus \cap \Omega_k\right) + \mathbb{P}\left(\Omega_k^c\right) \\ &\leq \frac{k_0}{\Theta} k^{2\left(\frac{1}{4}-\beta\right)-2\varepsilon-2\theta_0} + \frac{k_0}{\Theta} k^{-2\varepsilon} \lambda_N^{-2\left(\frac{1}{4}-\beta\right)+2\theta_1} + \frac{\tilde{C}_5}{\log k^{-\varepsilon}}. \end{aligned}$$

Letting  $N \nearrow \infty$ , then  $k \searrow 0$  and finally  $\Theta \nearrow \infty$  in the last line we easily conclude the proof of the corollary.  $\square$

To close this section let us make some few remarks. Instead of the scheme (25) we could also use a fully-implicit scheme. More precisely, for any  $j \in \{0, \dots, M-1\}$  we look for a  $\mathcal{F}_{t_j}$ -measurable random variable  $\mathcal{U}^j \in \mathbb{H}_N$  such that for any  $w \in \mathbb{V}$

$$(33) \quad \begin{cases} \mathcal{U}^0 = \pi_N \mathbf{u}_0, \\ \langle \mathcal{U}^{j+1} - \mathcal{U}^j + k[\pi_N A \mathcal{U}^{j+1} + \pi_N B(\mathcal{U}^{j+1}, \mathcal{U}^{j+1})], w \rangle = \langle w, \pi_N G(\mathcal{U}^j) \Delta_{j+1} W \rangle, \end{cases}$$

where  $\Delta_{j+1} W := W(t_{j+1}) - W(t_j)$ ,  $j \in \{0, \dots, M-1\}$ . We have the following theorem:

**Theorem 2.11.** *Let the assumptions (B1)-(B3) and (G) hold and assume that  $\mathbf{u}_0 \in L^{16}(\Omega; \mathbb{H}) \cap L^8(\Omega; \mathbb{V}_{\frac{1}{4}})$ . Let  $N$  and  $M$  be two fixed positive integers. Then,*

- (a) *for any  $j \in \{0, \dots, M-1\}$  there exists a unique  $\mathcal{F}_{t_j}$ -measurable random variable  $\mathcal{U}^j \in \mathbb{H}_N$  satisfying (33) and the estimates (26) and (30).*
- (b) *For any  $\beta \in [0, \frac{1}{4})$  there exists a constant  $k_0 > 0$  such that for any small number  $\varepsilon > 0$  we have*

$$(34) \quad \max_{1 \leq j \leq M} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \|\mathbf{u}(t_j) - \mathcal{U}^j\|_\beta^2 \right) + 2k \mathbb{E} \left( \mathbf{1}_{\Omega_k} \sum_{j=1}^M \|\mathbf{u}(t_j) - \mathcal{U}^j\|_{\frac{1}{2}+\beta}^2 \right) < k_0 k^{-2\varepsilon} [k^{2\left(\frac{1}{4}-\beta\right)} + \lambda_N^{-2\left(\frac{1}{4}-\beta\right)}],$$

where

$$\Omega_k = \left\{ \omega : \sup_{t \in [0, T]} \|\mathbf{u}(t, \omega)\|_{\frac{1}{4}}^2 < \log k^{-\varepsilon}, \max_{0 \leq j \leq M} \|\mathcal{U}^j(\omega)\|_{\frac{1}{4}}^2 < \log k^{-\varepsilon} \right\}.$$

- (c) *Moreover, for any small number  $\varepsilon > 0$ , any  $\theta_0 \in (0, \frac{1}{4} - \beta - \varepsilon)$  and  $\theta_1 \in (0, \frac{1}{4} - \beta)$*

$$(35) \quad \lim_{\Theta \nearrow \infty} \lim_{k \searrow 0} \lim_{N \nearrow \infty} \max_{1 \leq j \leq M} \mathbb{P} \left( \|\mathbf{u}(t_j) - \mathcal{U}^j\|_\beta^2 + k^{\frac{1}{2}} \left( \sum_{j=1}^M \|\mathbf{u}(t_j) - \mathcal{U}^j\|_{\frac{1}{2}+\beta}^2 \right)^{\frac{1}{2}} \geq \Theta [k^{\theta_0} + \lambda_N^{-\theta_1}] \right) = 0.$$

*Proof.* The arguments for the proof of this theorem are very similar to those of the proofs of Proposition 2.7, Theorem 2.8 and Corollary 2.10, thus we omit them.  $\square$

### 3. EXISTENCE AND STABILITY ANALYSIS OF THE SCHEME: PROOF OF PROPOSITION 2.7

In this section we will show that for any  $j \in \{0, \dots, M-1\}$  the numerical scheme (25) admits at least one solution  $\mathcal{U}^j \in \mathbb{H}_N$ . We will also show that (25) is stable in  $D(A^{\frac{1}{4}})$ , see Proposition 2.7 for more precision.

*Proof of Proposition 2.7.* As we mentioned in Subsection 2.3 we will only prove the existence, measurability and the estimates (29) and (30). The proof of the inequalities (26)-(28) will be omitted because it is very similar to the proof of [17] (see also [11]).

*Proof of the existence.* We first establish that for any  $j \in \{0, \dots, M-1\}$  there exists  $\mathcal{U}^j \in \mathbb{H}_N$  satisfying the numerical scheme (25). To this end, let us fix  $\omega \in \Omega$  and for a given  $\mathcal{U}^j \in \mathbb{H}_N$  consider the map  $\Lambda_\omega^j : \mathbb{H}_N \rightarrow \mathbb{H}_N$  defined by

$$\langle \Lambda_\omega^j(\mathbf{v}), \psi \rangle = \langle \mathbf{v} - \mathcal{U}^j(\omega), \psi \rangle + k \langle A\mathbf{v} + \pi_N B(\mathcal{U}^j(\omega), \mathbf{v}), \psi \rangle - \langle \psi, \pi_N G(\mathcal{U}^j(\omega)) \Delta_{j+1} W(\omega) \rangle$$

for any  $\psi \in H_N$ . Note that since  $H_N \subset D(A)$  the map  $\Lambda_\omega^j$  is well-defined. From assumptions **(B1)** and **(G)** and the linearity of  $A$  it is clear that for given  $U^j$  the map  $\Lambda_\omega^j$  is continuous. Furthermore, using Hölder's inequality, the fact that  $\lambda_1 |\psi|^2 \leq \|\psi\|_{\frac{1}{2}}^2$ ,  $\psi \in V$  and assumptions **(B2)** and **(G)** we derive that

$$\begin{aligned} \langle \Lambda_\omega^j \mathbf{v}, \mathbf{v} \rangle &\geq |\mathbf{v}|^2 \left( \lambda_1 k + \frac{1}{2} - \frac{k}{2} \right) - \frac{|U^j(\omega)|^2}{2} (1 + \|\Delta_{j+1} W(\omega)\|_{\mathcal{H}}^2 C_2^2) - \frac{1}{2} \|\Delta_{j+1} W(\omega)\|_{\mathcal{H}}^2 C_2^2 \\ &\geq \gamma |\mathbf{v}|^2 - \Gamma_\omega^j. \end{aligned}$$

Since  $k < 1$ , and by Assumption **(N)**,  $\|\Delta_{j+1} W\|_{\mathcal{H}}^2 < \infty$ , the constant  $\gamma$  is positive and  $\mu_j = \sqrt{\frac{\Gamma_\omega^j}{\gamma}} < \infty$  whenever  $|U^j|^2 < \infty$ . Thus, we have  $\langle \Lambda_\omega^j \mathbf{v}, \mathbf{v} \rangle \geq 0$  for any  $\mathbf{v} \in \mathcal{H}_N^j(\omega) := \{\psi \in H_N; |\psi| = R\mu_j\}$  where  $R > 1$  is an arbitrary constant. Since  $U^0 = \pi_N \mathbf{u}_0$  is given, we can conclude from the above observations and Brouwer fixed point theorem that there exists at least one  $U^1 \in H_N$  satisfying

$$\Lambda_\omega^0(U^1) = 0 \text{ and } |U^1| \leq R\mu_0.$$

In a similar way, assuming that  $U^j \in H_N$ , we infer that there exists at least one  $U^{j+1} \in H_N$  such that

$$\Lambda_\omega^j(U^{j+1}) = 0 \text{ and } |U^{j+1}| \leq R\mu_j.$$

Therefore, we have to prove by induction that given  $U^0 \in H_N$  and a  $\mathcal{H}$ -valued Wiener process  $W$ , for each  $j$ , there exists a sequence  $\{U^j; j = 1, \dots, M\} \subset H_N$  satisfying the algorithm (25).

*Proof of the measurability.* In order to prove the  $\mathcal{F}_{t_j}$ -measurability of  $U^j$  it is sufficient to show that for each  $j \in \{1, \dots, M\}$  one can find a Borel measurable map  $\mathcal{E}_j : H_N \times \mathcal{H} \rightarrow H_N$  such that  $U^j = \mathcal{E}_j(U^{j-1}, \Delta_j W)$ . In fact, if such claim is true then by exploiting the  $\mathcal{F}_{t_j}$ -measurability of  $\Delta_j W$  one can argue by induction and show that if  $U^0$  is  $\mathcal{F}_0$ -measurable then  $\mathcal{E}_j(U^{j-1}, \Delta_j W)$  is  $\mathcal{F}_{t_j}$ -measurable, hence  $U^j$  is  $\mathcal{F}_{t_j}$ -measurable. Thus, it remains to prove the existence of  $\mathcal{E}_j$ . For this purpose we will closely follow [23]. Let  $\mathcal{P}(H_N)$  be the set of subsets of  $H_N$  and consider a multivalued map  $\mathcal{E}_{j+1}^S : H_N \times \mathcal{H} \rightarrow \mathcal{P}(H_N)$  such that for each  $(U^j, \eta_{j+1})$ ,  $\mathcal{E}_{j+1}^S(U^j, \eta_{j+1})$  denotes the set of solutions  $U^{j+1}$  of (25). From the existence result above we deduce that  $\mathcal{E}_{j+1}^S$  maps  $H_N \times \mathcal{H}$  to nonempty closed subsets of  $H_N$ . Furthermore, since we are in the finite dimensional space  $H_N$ , we can prove, by using the assumptions **(B1)** and **(G)** and the sequential characterization of the closed graph theorem, that the graph of  $\mathcal{E}_{j+1}^S$  is closed. From these last two facts and [4, Theorem 3.1] we can find a univocal map  $\mathcal{E}_{j+1} : H_N \times \mathcal{H} \rightarrow H_N$  such that  $\mathcal{E}_j(U^j, \eta_{j+1}) \in \mathcal{E}_{j+1}^S(U^j, \eta_{j+1})$  and  $\mathcal{E}_j$  is measurable when  $H_N \times \mathcal{H}$  and  $H_N$  are equipped with their respective Borel  $\sigma$ -algebra. This completes the proof of the measurability of the solutions of (25).

*Proof of (26)-(28).* Thanks to the assumption **(B2)**, the proof of the inequalities (26)-(28) is very similar to the proof of [17], so we omit it and we directly proceed to the proof of the estimates (29) and (30).

*Proof of (29).* Taking  $w = 2A^{\frac{1}{2}}U^{j+1}$  in (25), using the Cauchy-Schwarz inequality and the identity

$$(36) \quad ((\mathbf{v} - \mathbf{x}, 2\mathbf{v})) = \|\mathbf{v}\|^2 - \|\mathbf{x}\|^2 + \|\mathbf{v} - \mathbf{x}\|^2, \quad (\mathbf{v}, \mathbf{x} \text{ are elements of a Hilbert space with norm } \|\cdot\|)$$

yield

$$\begin{aligned} &\|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 + \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + 2k\|U^{j+1}\|_{\frac{3}{4}}^2 \\ &\leq 2k\|\pi_N B(U^j, U^{j+1})\| \|U^{j+1}\|_{\frac{1}{2}} + 2\|\pi_N G(U^j) \Delta_{j+1} W\|_{\frac{1}{4}} \|U^{j+1} - U^j\|_{\frac{1}{4}} \\ &\quad + 2\langle A^{\frac{1}{4}}U^j, A^{\frac{1}{4}}\pi_N G(U^j) \Delta_{j+1} W \rangle. \end{aligned}$$

Using the fact that  $\|\pi_N\|_{\mathcal{L}(H, H_N)} \leq 1$ , we obtain

$$(37) \quad \begin{aligned} & \|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 + \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + 2k\|U^{j+1}\|_{\frac{3}{4}}^2 \\ & \leq 2k|\mathbf{B}(U^j, U^{j+1})| \|U^{j+1}\|_{\frac{1}{2}} + 2\|G(U^j)\Delta_{j+1}W\|_{\frac{1}{4}} \|U^{j+1} - U^j\|_{\frac{1}{4}} \\ & \quad + 2\langle A^{\frac{1}{4}}U^j, A^{\frac{1}{4}}\pi_N G(U^j)\Delta_{j+1}W \rangle. \end{aligned}$$

Using Assumption **(B1)'**, the complex interpolation inequality in [43, Theorem 1.9.3, pp 59], the Young inequality, and the continuous embedding  $V_{\frac{1}{2}} \subset V_{\frac{1}{4}}$  we obtain

$$(38) \quad \begin{aligned} 2|\mathbf{B}(U^j, U^{j+1})| \|U^{j+1}\|_{\frac{1}{2}} & \leq C|U^j|^4 \|U^{j+1}\|_{\frac{1}{4}}^2 + \|U^{j+1}\|_{\frac{3}{4}}^2 \\ & \leq C|U^j|^4 \|U^{j+1}\|_{\frac{1}{2}}^2 + \|U^{j+1}\|_{\frac{3}{4}}^2, \end{aligned}$$

which implies that

$$(39) \quad \begin{aligned} \|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 + \frac{1}{2}\|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + 2k\|U^{j+1}\|_{\frac{3}{4}}^2 & \leq 2Ck|U^j|^4 \|U^{j+1}\|_{\frac{1}{2}}^2 + 4\|G(U^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2 \\ & \quad + 2\langle A^{\frac{1}{4}}U^j, A^{\frac{1}{4}}\pi_N G(U^j)\Delta_{j+1}W \rangle. \end{aligned}$$

Since  $U^j$  is a constant, adapted and hence progressively measurable process, it is not difficult to prove that

$$2\mathbb{E}\langle A^{\frac{1}{4}}U^j, A^{\frac{1}{4}}\pi_N G(U^j)\Delta_{j+1}W \rangle = 0.$$

Using (27) and (28) with  $p = 2$  and  $p = 3$  respectively, we easily prove that there exists a constant  $C > 0$ , depending only on  $T$ , such that

$$(40) \quad k\mathbb{E}\left(\sum_{j=0}^{M-1} |U^j|^4 \|U^{j+1}\|_{\frac{1}{2}}^2\right) \leq \left(\mathbb{E}\max_{1 \leq m \leq M} |U^m|^8\right)^{\frac{1}{2}} \left(\mathbb{E}\left(k\sum_{j=1}^M \|U^j\|_{\frac{1}{2}}^2\right)^2\right)^{\frac{1}{2}} \leq C(1 + \mathbb{E}|u_0|^8)^2.$$

Now, since  $U^j$  is  $\mathcal{F}_j$ -measurable and  $\Delta_{j+1}W$  is independent of  $\mathcal{F}_j$ , we infer that there exists a constant  $C > 0$  such that for any  $j \in \{0, \dots, M-1\}$

$$(41) \quad \begin{aligned} \mathbb{E}\left(\|G(U^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2\right) & \leq \mathbb{E}\left(\mathbb{E}\left(\|G(U^j)\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 \|\Delta_{j+1}W\|_{\mathcal{H}}^2 \mid \mathcal{F}_j\right)\right) \\ & = m\mathbb{E}\left(\mathbb{E}\left(\|G(U^j)\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 \mathbb{E}\left(\|\Delta_{j+1}W\|_{\mathcal{H}}^2 \mid \mathcal{F}_j\right)\right)\right) \\ & \leq Ck(\operatorname{tr}Q)^{\frac{1}{2}}(1 + \mathbb{E}\|U^j\|_{\frac{1}{4}}^2), \end{aligned}$$

where (8) and Assumption **(G)** along with Remark 2.1-(b) were used to derive the last line of the above chain of inequalities.

Now taking the mathematical expectation in (39), summing both sides of the resulting equations from  $j = 0$  to  $m-1$  and using the last three observations imply

$$\begin{aligned} & \max_{1 \leq m \leq M} \mathbb{E}\|U^m\|_{\frac{1}{4}}^2 + \frac{1}{2}\mathbb{E}\left(\sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2\right) + 2k\mathbb{E}\sum_{j=1}^M \|U^j\|_{\frac{3}{4}}^2 \\ & \leq CT + \mathbb{E}\|u_0\|_{\frac{1}{4}}^2 + C\operatorname{Tr}Qk\sum_{m=1}^M \max_{1 \leq j \leq m} \mathbb{E}\|U^j\|_{\frac{1}{4}}^2, \end{aligned}$$

from which along with the discrete Gronwall lemma we infer that there exists a constant  $C > 0$  such that

$$(42) \quad \max_{1 \leq m \leq M} \mathbb{E}\|U^m\|_{\frac{1}{4}}^2 + \frac{1}{2}\mathbb{E}\left(\sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2\right) + 2k\mathbb{E}\sum_{j=1}^M \|U^j\|_{\frac{3}{4}}^2 \leq C(1 + \mathbb{E}\|u_0\|_{\frac{1}{4}}^2 + [\mathbb{E}|u_0|^8]^2).$$

Note that from (39) we can derive that there exists a constant  $C > 0$  such that

$$\begin{aligned} \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^2 &\leq \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^2 + Ck \mathbb{E} \sum_{j=0}^{M-1} |U^j|^4 \|U^{j+1}\|_{\frac{1}{2}}^2 + \mathbb{E} \sum_{j=0}^{M-1} \|G(U^j) \Delta_{j+1} W\|_{\frac{1}{4}}^2 \\ &\quad + 2\mathbb{E} \max_{1 \leq m \leq M} \sum_{j=0}^{m-1} \langle A^{\frac{1}{4}} \pi_N G(U^j) \Delta_{j+1} W, A^{\frac{1}{4}} U^j \rangle \\ &=: \sum_{i=1}^4 I_i. \end{aligned}$$

Arguing as in [11, proof of (3.9)] we can establish that

$$I_4 \leq \frac{1}{2} \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^2 + \frac{1}{2} \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^2 + Ck \sum_{j=0}^{M-1} \mathbb{E} \|U^j\|_{\frac{1}{4}}^2,$$

which altogether with (42) yields that

$$I_4 \leq \frac{1}{2} \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^2 + C(1 + \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^2).$$

Using the same idea as in the proof of (41) and using (42) we infer that

$$I_3 \leq C(1 + \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^2).$$

Using these two estimates and the inequality (40) we derive that there exists a constant  $C > 0$  such that

$$\mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^2 \leq C(1 + \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^2 + [\mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^8]^2),$$

which along with (42) completes the proof of (29).

Now, we continue with the derivation of an estimate of  $\max_{1 \leq m \leq M} \mathbb{E} \|U^m\|_{\frac{1}{4}}^4$ . Multiplying (37) by  $\|U^{j+1}\|_{\frac{1}{4}}^2$  and using identity (36) and then summing both sides of the resulting equation from  $j = 0$  to  $m - 1$  implies

$$\begin{aligned} (43) \quad &\frac{1}{2} \|U^m\|_{\frac{1}{4}}^4 + \frac{1}{2} \sum_{j=0}^{m-1} \left| \|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 \right|^2 + \sum_{j=0}^{m-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + 2k \sum_{j=0}^{m-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \|U^{j+1}\|_{\frac{3}{4}}^2 \\ &\leq \frac{1}{2} \|\mathbf{u}_0\|_{\frac{1}{4}}^4 + Ck \sum_{j=0}^{m-1} |B(U^j, U^{j+1})|^2 \|U^{j+1}\|_{\frac{1}{2}}^2 \|U^{j+1}\|_{\frac{1}{4}}^2 \\ &\quad + 2 \sum_{j=0}^{m-1} \langle A^{\frac{1}{4}} [U^{j+1} - U^j], A^{\frac{1}{4}} \pi_N G(U^j) \Delta_{j+1} W \rangle \|U^{j+1}\|_{\frac{1}{4}}^2 \\ &\quad + 2 \sum_{j=0}^{m-1} \langle A^{\frac{1}{4}} U^j, A^{\frac{1}{4}} \pi_N G(U^j) \Delta_{j+1} W \rangle \|U^{j+1}\|_{\frac{1}{4}}^2 \\ &=: \frac{1}{2} \|\mathbf{u}_0\|_{\frac{1}{4}}^4 + J_1 + J_2 + J_3. \end{aligned}$$

Thanks to the estimate (38) we can estimate  $J_1$  as follows

$$\mathbb{E} J_1 \leq CK \mathbb{E} \sum_{j=0}^{M-1} |U^j|^4 \|U^{j+1}\|_{\frac{1}{4}}^4 + k \mathbb{E} \sum_{j=0}^{M-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \|U^{j+1}\|_{\frac{3}{4}}^2 =: J_{1,1} + J_{1,2}.$$

Since the second term  $J_{1,2}$  can be absorbed in the LHS later on, we will focus on estimating the second term  $J_{1,1}$ . We have

$$\begin{aligned}
 J_{1,1} &\leq Ck \sum_{j=0}^{M-1} |\mathbf{U}^j|^4 |\mathbf{U}^{j+1}|^2 \|\mathbf{U}^{j+1}\|_{\frac{1}{2}}^2 \\
 &\leq C \left( \mathbb{E} \max_{0 \leq j \leq M-1} [|\mathbf{U}^j|^8 |\mathbf{U}^{j+1}|^4] \right)^{\frac{1}{2}} \left( \mathbb{E} \left[ k \sum_{j=1}^M \|\mathbf{U}^j\|_{\frac{1}{2}}^2 \right]^2 \right)^{\frac{1}{2}} \\
 &\leq C \left( \mathbb{E} \left[ \max_{0 \leq j \leq M-1} |\mathbf{U}^j|^{12} \right] \right)^{\frac{1}{2}} \left( \mathbb{E} \left[ k \sum_{j=1}^M \|\mathbf{U}^j\|_{\frac{1}{2}}^2 \right]^2 \right)^{\frac{1}{2}} \\
 &\leq C(1 + \mathbb{E}|\mathbf{u}_0|^{16}),
 \end{aligned}$$

where (27) and (28) are used to obtain the last line. Hence,

$$\mathbb{E}J_1 \leq C(1 + \mathbb{E}|\mathbf{u}_0|^{16}) + \mathbb{E}k \sum_{j=0}^{M-1} \left( \|\mathbf{U}^{j+1}\|^2 - \frac{1}{4} \|\mathbf{U}^{j+1}\|_{\frac{3}{4}}^2 \right).$$

Now we estimate  $J_2$  as follows

$$\begin{aligned}
 \mathbb{E}J_2 &\leq C\mathbb{E} \sum_{j=0}^{M-1} \|G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2 \left( \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 + \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right) + \frac{1}{2}\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{U}^{j+1} - \mathbf{U}^j\|_{\frac{1}{4}}^2 \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 \\
 &\leq C\mathbb{E} \sum_{j=0}^{M-1} \|G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^4 + C\mathbb{E} \sum_{j=0}^{M-1} \|G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2 \|\mathbf{U}^j\|_{\frac{1}{4}}^2 + \frac{1}{8}\mathbb{E} \sum_{j=0}^{M-1} \left| \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right|^2 \\
 &\quad + \frac{1}{2}\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{U}^{j+1} - \mathbf{U}^j\|_{\frac{1}{4}}^2 \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2.
 \end{aligned}$$

As long as  $J_3$  is concerned we have

$$\begin{aligned}
 \mathbb{E}J_3 &= 2\mathbb{E} \sum_{j=0}^{m-1} \langle \mathbf{A}^{\frac{1}{4}}\mathbf{U}^j, \mathbf{A}^{\frac{1}{4}}\pi_N G(\mathbf{U}^j)\Delta_{j+1}W \rangle \|\mathbf{U}^j\|_{\frac{1}{4}}^2 + 2\mathbb{E} \sum_{j=0}^{m-1} \langle \mathbf{A}^{\frac{1}{4}}\mathbf{U}^j, \mathbf{A}^{\frac{1}{4}}G(\mathbf{U}^j)\Delta_{j+1}W \rangle \left( \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right) \\
 &= 2\mathbb{E} \sum_{j=0}^{m-1} \langle \mathbf{A}^{\frac{1}{4}}\mathbf{U}^j, \mathbf{A}^{\frac{1}{4}}\pi_N G(\mathbf{U}^j)\Delta_{j+1}W \rangle \left( \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right) \\
 &\leq C\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{A}^{\frac{1}{4}}G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2 \|\mathbf{U}^j\|_{\frac{1}{4}}^2 + \frac{1}{8}\mathbb{E} \sum_{j=0}^{M-1} \left| \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right|^2
 \end{aligned}$$

because for any  $j$

$$\mathbb{E} \langle \mathbf{A}^{\frac{1}{4}}\mathbf{U}^j, \mathbf{A}^{\frac{1}{4}}\pi_N G(\mathbf{U}^j)\Delta_{j+1}W \rangle \|\mathbf{U}^j\|_{\frac{1}{4}}^2 = 0.$$

By a similar idea as used to derive (41) we can prove that

$$C\mathbb{E} \sum_{j=0}^{M-1} \|G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^4 + C\mathbb{E} \sum_{j=0}^{M-1} \|G(\mathbf{U}^j)\Delta_{j+1}W\|_{\frac{1}{4}}^2 \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \leq C + Ck\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{U}^j\|_{\frac{1}{4}}^4.$$

Thus,

$$\mathbb{E}[J_2 + J_3] \leq C + Ck\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{U}^j\|_{\frac{1}{4}}^4 + \frac{1}{4}\mathbb{E} \sum_{j=0}^{M-1} \left| \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2 - \|\mathbf{U}^j\|_{\frac{1}{4}}^2 \right|^2 + \frac{1}{2}\mathbb{E} \sum_{j=0}^{M-1} \|\mathbf{U}^{j+1} - \mathbf{U}^j\|_{\frac{1}{4}}^2 \|\mathbf{U}^{j+1}\|_{\frac{1}{4}}^2.$$

Taking the mathematical expectation in (43) and by plugging the information about  $J_i$ ,  $i = 1, 2, 3$  in the resulting equation yield

$$\begin{aligned} & \max_{1 \leq m \leq M} \frac{1}{2} \mathbb{E} \|U^m\|_{\frac{1}{4}}^4 + \frac{1}{4} \mathbb{E} \sum_{j=0}^{M-1} \left| \|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 \right|^2 \\ & + \frac{1}{2} \mathbb{E} \sum_{j=0}^{M-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 + k \mathbb{E} \sum_{j=1}^M \|U^j\|_{\frac{1}{4}}^2 \|U^j\|_{\frac{3}{4}}^2 \\ & \leq C(1 + \mathbb{E} |\mathbf{u}_0|^{12} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4) + Ck \mathbb{E} \sum_{j=0}^{M-1} \|U^j\|_{\frac{1}{4}}^4, \end{aligned}$$

which along with the Gronwall inequality yields

$$\max_{1 \leq m \leq M} \frac{1}{2} \mathbb{E} \|U^m\|_{\frac{1}{4}}^4 \leq C(1 + \mathbb{E} |\mathbf{u}_0|^{12} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4).$$

The latter inequality is used in the former one to derive that

$$(44) \quad \begin{aligned} & \max_{1 \leq m \leq M} \frac{1}{2} \mathbb{E} \|U^m\|_{\frac{1}{4}}^4 + \frac{1}{4} \mathbb{E} \sum_{j=0}^{M-1} \left| \|U^{j+1}\|_{\frac{1}{4}}^2 - \|U^j\|_{\frac{1}{4}}^2 \right|^2 + \frac{1}{2} \mathbb{E} \sum_{j=0}^{M-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 \\ & + k \mathbb{E} \sum_{j=1}^M \|U^j\|_{\frac{1}{4}}^2 \|U^j\|_{\frac{3}{4}}^2 \leq C(1 + \mathbb{E} |\mathbf{u}_0|^{12} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4). \end{aligned}$$

Now we continue our analysis with the estimation of  $\mathbb{E} \max_{1 \leq j \leq M} \|U^j\|_{\frac{1}{4}}^4$ . To start with this analysis, we easily derive from (43) the following inequality

$$\begin{aligned} \max_{1 \leq m \leq M} \frac{1}{2} \mathbb{E} \|U^m\|_{\frac{1}{4}}^4 & \leq Ck \sum_{j=0}^{M-1} \|U^j\|^4 \|U^{j+1}\|^2 \|U^{j+1}\|_{\frac{1}{2}}^2 \\ & + C \sum_{j=0}^{M-1} \left( \|G(U^j) \Delta_{j+1}\|_{\frac{1}{4}}^4 + \|G(U^j) \Delta_{j+1}\|_{\frac{1}{4}}^2 \|U^j\|_{\frac{1}{4}}^2 \right) \\ & + \max_{0 \leq j \leq M-1} \sum_{\ell=0}^{j-1} \langle A^{\frac{1}{4}} U^\ell, A^{\frac{1}{4}} \pi_N G(U^\ell) \Delta_{\ell+1} W \rangle \|U^\ell\|_{\frac{1}{4}}^2 =: J_1 + J_2 + J_3. \end{aligned}$$

Arguing as in the proof of (41) and using (44), the mathematical expectation of  $J_1 + J_2$  can be estimated as follows

$$\mathbb{E}(J_1 + J_2) \leq C \mathbb{E}(1 + |\mathbf{u}_0|^{16} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4).$$

The same idea as used in the proof of [11, inequality (3.15)] yields

$$\mathbb{E} J_3 \leq \frac{1}{4} \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^4 + C \mathbb{E} \|\mathbf{u}_0\|_{\frac{1}{4}}^4 + Ck \mathbb{E} \sum_{j=0}^{M-1} \|U^j\|_{\frac{1}{4}}^4,$$

from which altogether with (44) we infer that

$$\mathbb{E} J_3 \leq C \mathbb{E}(1 + |\mathbf{u}_0|^{16} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4) + \frac{1}{4} \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^4.$$

Thus, summing up we have shown that there exists a constant  $C > 0$  such that

$$(45) \quad \mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^4 \leq C \mathbb{E}(1 + |\mathbf{u}_0|^{16} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4).$$



Now, we estimate  $\mathbb{E} \left( \sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 \right)^2 + \mathbb{E} \left( k \sum_{j=1}^M \|U^j\|_{\frac{3}{4}}^2 \right)^2$ . To do this we first observe that from (39) we infer that

$$(46) \quad \left( \frac{1}{2} \sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 \right)^2 + \left( 2k \sum_{j=0}^{M-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \right)^2 \leq C \left( k \sum_{j=0}^{M-1} |U^j|^4 \|U^{j+1}\|_{\frac{1}{2}}^2 \right)^2 \\ + C \left( \sum_{j=0}^{M-1} \|G(U^j) \Delta_{j+1} W\|_{\frac{1}{4}}^2 \right)^2 + C \left( \sum_{j=0}^{M-1} \langle A^{\frac{1}{4}} U^j, A^{\frac{1}{4}} \pi_N G(U^j) \Delta_{j+1} W \rangle \right)^2.$$

Then, using the same strategies to estimate the  $J_i$ -s (or  $J_j$ ), the sum of the three terms in the right hand side of the above quality can be bounded from above by

$$\left[ \mathbb{E} \left( \max_{0 \leq j \leq M} |U^j|^{16} \right) \right]^{\frac{1}{2}} \left[ \mathbb{E} \left( k \sum_{j=1}^M \|U^j\|_{\frac{1}{2}}^2 \right)^4 \right]^{\frac{1}{2}} + CMk^2 \sum_{j=0}^M \mathbb{E} \|U^j\|_{\frac{1}{4}}^4 + Ck \sum_{j=0}^M \mathbb{E} \|U^j\|_{\frac{1}{4}}^4,$$

which along with the estimate for  $\mathbb{E} \max_{1 \leq m \leq M} \|U^m\|_{\frac{1}{4}}^4$  and the inequalities (27) and (28) implies that

$$(47) \quad \left( \frac{1}{2} \sum_{j=0}^{M-1} \|U^{j+1} - U^j\|_{\frac{1}{4}}^2 \right)^2 + \left( 2k \sum_{j=0}^{M-1} \|U^{j+1}\|_{\frac{1}{4}}^2 \right)^2 \leq \mathbb{E}(1 + |\mathbf{u}_0|^{16} + \|\mathbf{u}_0\|_{\frac{1}{4}}^4).$$

The last estimate along with (45) completes the proof of (30) and hence the whole proposition.  $\square$

#### 4. ERROR ANALYSIS OF THE NUMERICAL SCHEME (25): PROOF OF THEOREM 2.8

This section is devoted to the analysis of the error  $e_j = \mathbf{u}(t_j) - U^j$  at the time  $t_j$  between the exact solution  $\mathbf{u}$  of (1) and the approximate solution given by (25). Since the precise statement of the convergence rate is already given in Theorem 2.8, we proceed directly to the promised proof of Theorem 2.8.

Before giving the proof of Theorem 2.8 we state and prove the following important result.

**Lemma 4.1.** *Let  $\beta$  be as in Theorem 2.8. Then,*

(i) *there exists a constant  $C_7 > 0$  such that*

$$(48) \quad \mathbb{E} \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\beta}^2 \leq C_7 [(t-s)^{2-2\beta} + (t-s)^{2(\frac{1}{4}-\beta)} + (t-s)],$$

*for any  $t, s \geq 0$  and  $t \neq s$ .*

(ii) *There also exists a positive constant  $C_8$  such that*

$$(49) \quad \mathbb{E} \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr \leq C_8 \left( (t-s)^{\frac{3}{2}-2\beta} + (t-s)^{2(\frac{1}{4}-\beta)} + (t-s)^{2-2\beta} \right),$$

*for any  $t > s \geq 0$ .*

*Proof of Lemma 4.1.* As in the statement of the lemma we divide the proof into two parts.

*Proof of item (i).* Let  $t, s \in [0, T]$  such that  $t \neq s$ . Without loss of generality we assume that  $t > s$ . Thanks to (12) of Remark 2.5 we have

$$\|\mathbf{u}(t) - \mathbf{u}(s)\|_{\beta}^2 \leq C |A^{\beta-\frac{1}{4}} (\mathbf{I} - e^{-(t-s)A}) A^{\frac{1}{4}} \mathbf{u}(s)|^2 + C \left| \int_s^t A^{\beta} e^{-(t-r)A} B(\mathbf{u}(r), \mathbf{u}(r)) dr \right|^2 \\ + C \left| \int_s^t A^{\beta} e^{-(t-r)A} G(\mathbf{u}(r)) dW(r) \right|^2.$$

Before proceeding further we recall that there exists a constant  $C > 0$  such that for any  $\gamma > 0$  and  $t \geq 0$ , we have

$$\|A^{-\gamma} (\mathbf{I} - e^{-tA})\|_{\mathcal{L}(\mathbf{H})} \leq Ct^{\gamma}.$$

Applying this inequality, the Hölder inequality, Assumption **(B1)'**, the Itô isometry and Assumption **(G)** imply

$$\begin{aligned} \mathbb{E} \left( \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\beta}^2 \right) &\leq C(t-s) \mathbb{E} \left( \int_s^t (t-r)^{-2\beta} \|\mathbf{u}(r)\|_{\frac{1}{4}}^2 \|\mathbf{u}(r)\|_{\frac{1}{2}-\frac{1}{4}}^2 dr \right) \\ &\quad + C(t-s)^{2(\frac{1}{4}-\beta)} \mathbb{E} \|\mathbf{u}(s)\|_{\frac{1}{4}}^2 + \mathbb{E} \int_s^t |e^{-(t-r)A} \mathbf{A}^{\beta} \mathbf{G}(\mathbf{u}(r))|^2 dr \\ &\leq C(t-s)^{2-2\beta} \mathbb{E} \left( \sup_{r \in [s,t]} \|\mathbf{u}(r)\|_{\frac{1}{4}}^2 \sup_{r \in [s,t]} \|\mathbf{u}(r)\|_{\frac{1}{2}-\frac{1}{4}}^2 \right) \\ &\quad + C[(t-s)^{2(\frac{1}{4}-\beta)} + (t-s)] \mathbb{E} \left( \sup_{r \in [s,t]} \|\mathbf{u}(r)\|_{\frac{1}{4}}^4 \right), \end{aligned}$$

from which along with (15) we easily infer that

$$\mathbb{E} \left( \|\mathbf{u}(t) - \mathbf{u}(s)\|_{\beta}^2 \right) \leq C[(t-s)^{2-2\beta} + (t-s)^{2(\frac{1}{4}-\beta)} + (t-s)].$$

Thus, we have just finished the proof of the first part of the lemma.

*Proof of item (ii).* Let  $t > s \geq 0$ . Using (12) of Remark 2.5, it is not difficult to see that

$$\begin{aligned} \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr &\leq C \int_s^t \left( \int_r^t |A^{\frac{1}{2}+\beta} e^{-(t-\tau)A} \mathbf{B}(\mathbf{u}(\tau), \mathbf{u}(\tau))| d\tau \right)^2 dr \\ &\quad + C \int_s^t \left| \int_r^t A^{\frac{1}{4}+\beta} e^{-(t-\tau)A} [A^{\frac{1}{4}} \mathbf{G}(\mathbf{u}(\tau))] dW(\tau) \right|^2 dr \\ &\quad + C \int_s^t |A^{\beta-\frac{1}{4}} (e^{-(t-r)A} - \mathbf{I}) A^{\frac{3}{4}} \mathbf{u}(s)|^2 dr, \end{aligned}$$

from which and the assumption on **B** we infer that

$$\begin{aligned} \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr &\leq C \sup_{0 \leq \tau \leq T} \left( \|\mathbf{u}(\tau)\|_{\frac{1}{4}}^2 \|\mathbf{u}(\tau)\|_{\frac{1}{2}-\frac{1}{4}}^2 \right) \int_s^t \left( \int_r^t (t-\tau)^{-\frac{1}{2}-\beta} d\tau \right)^2 dr \\ &\quad + C \int_s^t \left| \int_r^t A^{\frac{1}{4}+\beta} e^{-(t-\tau)A} [A^{\frac{1}{4}} \mathbf{G}(\mathbf{u}(\tau))] dW(\tau) \right|^2 dr \\ &\quad + C \int_s^t (t-r)^{2(\frac{1}{4}-\beta)} \|\mathbf{u}(s)\|_{\frac{3}{4}}^2 dr. \end{aligned}$$

Taking the mathematical expectation and using (15) yield

$$\begin{aligned} \mathbb{E} \left( \mathbb{1}_{\Omega_k} \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr \right) &\leq C(t-s)^{2-2\beta} + C(t-s)^{2(\frac{1}{4}-\beta)} \mathbb{E} \int_0^T \|\mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr \\ &\quad + \int_s^t \mathbb{E} \left( \left| \int_r^t A^{\frac{1}{4}+\beta} e^{-(t-\tau)A} A^{\frac{1}{4}} \mathbf{G}(\mathbf{u}(\tau)) dW(\tau) \right|^2 \right) dr. \end{aligned}$$

Owing to the Itô isometry, the assumption **(G)** and (15), we obtain

$$\begin{aligned} \mathbb{E} \left( \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr \right) &\leq \mathbb{E} \left( \sup_{0 \leq \tau \leq T} (1 + \|\mathbf{u}(\tau)\|_{\frac{1}{4}}^2) \right) \int_s^t \int_r^t (t-\tau)^{-\frac{1}{2}-2\beta} d\tau dr \\ &\quad + (t-s)^{2-2\beta} + (t-s)^{2(\frac{1}{4}-\beta)}, \end{aligned}$$

from which altogether with (15) we infer that there exists a constant  $C > 0$  such that

$$\mathbb{E} \left( \int_s^t \|\mathbf{u}(t) - \mathbf{u}(r)\|_{\frac{1}{2}+\beta}^2 dr \right) \leq C(t-s)^{2-2\beta} + C(t-s)^{2(\frac{1}{4}-\beta)} + C(t-s)^{\frac{3}{2}-2\beta},$$

for any  $t > s \geq 0$ . □

We now give the promised proof of Theorem 2.8.

*Proof of Theorem 2.8.* Since the embedding  $V_\beta \subset H$  is continuous for any  $\beta \in (0, \frac{1}{4})$ , it is sufficient to prove the main theorem for  $\beta \in (0, \frac{1}{4})$ .

Note that the numerical scheme (25) is equivalent to

$$(50) \quad (U^{j+1}, w) + \int_{t_j}^{t_{j+1}} \langle AU^{j+1} + \pi_N B(U^j, U^{j+1}), w \rangle ds = (U^j, w) + \int_{t_j}^{t_{j+1}} \langle w, \pi_N G(U^j) dW(s) \rangle$$

for any  $j \in \{1, \dots, M\}$  and  $w \in V$ . Integrating (1) and subtracting the resulting equation and the identity (50) term by term yield

$$(51) \quad (\mathbf{e}^{j+1} - \mathbf{e}^j, w) + \int_{t_j}^{t_{j+1}} \langle A\mathbf{e}^{j+1} + A(\mathbf{u}(s) - \mathbf{u}(t_{j+1})) + B(\mathbf{u}(s), \mathbf{u}(s)) - \pi_N B(U^j, U^{j+1}), w \rangle ds \\ = \int_{t_j}^{t_{j+1}} \langle w, [G(\mathbf{u}(s)) - \pi_N G(U^j)] dW(s) \rangle.$$

Observe that if  $\mathbf{v} \in D(A^{\frac{1}{2}+\alpha})$  with  $\alpha > \beta$ , then  $A^{2\beta}\mathbf{v} \in D(A^{\frac{1}{2}+\alpha-\beta}) \subset D(A^{\frac{1}{2}-\alpha})$ ,  $A\mathbf{v} \in D(A^{\alpha-\frac{1}{2}})$  and the duality product  $\langle A\mathbf{v}, A^{2\beta}\mathbf{v} \rangle$  is meaningful. Thus, we are permitted to take  $w = 2A^{2\beta}\mathbf{e}^{j+1}$  in (51) and derive that

$$\|\mathbf{e}^{j+1}\|_\beta^2 - \|\mathbf{e}^j\|_\beta^2 + \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 + 2k\|\mathbf{e}^{j+1}\|_{\frac{1}{2}+\beta}^2 - 2 \int_{t_j}^{t_{j+1}} \|A^{\frac{1}{2}+\beta}(\mathbf{u}(s) - \mathbf{u}(t_{j+1}))\|_{\frac{1}{2}+\beta} \|\mathbf{e}^{j+1}\|_{\frac{1}{2}+\beta} ds \\ \leq 2 \int_{t_j}^{t_{j+1}} \left| (A^{\beta-\frac{1}{2}}[B(\mathbf{u}(s), \mathbf{u}(s)) - \pi_N B(U^j, U^{j+1})], A^{\frac{1}{2}+\beta}\mathbf{e}^{j+1}) \right| ds \\ + 2 \int_{t_j}^{t_{j+1}} \langle A^{2\beta}\mathbf{e}^{j+1}, [G(\mathbf{u}(s)) - \pi_N G(U^j)] dW(s) \rangle,$$

where we have used the identity  $(\mathbf{v} - \mathbf{x}, 2A^{2\beta}\mathbf{v}) = \|\mathbf{v}\|_\beta^2 - \|\mathbf{x}\|_\beta^2 + \|\mathbf{v} - \mathbf{x}\|_\beta^2$ . Now, by using the identity  $\mathbf{v} = (\pi_N + [I - \pi_N])\mathbf{v}$ , the fact that

$$B(\mathbf{u}(s), \mathbf{u}(s)) - \pi_N B(U^j, U^{j+1}) = B(\mathbf{u}(s), \mathbf{u}(s)) - \pi_N B(\mathbf{u}(t_j), \mathbf{u}(t_{j+1})) \\ + \pi_N B(\mathbf{u}(t_j), \mathbf{u}(t_{j+1})) - B(U^j, U^{j+1}),$$

the Cauchy-Schwarz inequality, the Cauchy inequality  $ab \leq \frac{a^2}{4} + b^2$ ,  $a, b > 0$  and Assumption (B1) we obtain

$$(52) \quad \|\mathbf{e}^{j+1}\|_\beta^2 - \|\mathbf{e}^j\|_\beta^2 + \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 + k\|\mathbf{e}^{j+1}\|_{\frac{1}{2}+\beta}^2 \leq 2\mathcal{L}_j + 16C_0^2 \sum_{i=1}^5 \mathcal{N}_{j,i} + 2\mathcal{W}_j,$$

where for each  $j \in \{0, \dots, M-1\}$  the symbols  $\mathcal{L}_j$ ,  $\mathcal{N}_{j,i}$ ,  $i = 1, \dots, 5$ , and  $\mathcal{W}_j$  are defined by

$$\mathcal{L}_j := \int_{t_j}^{t_{j+1}} \|\mathbf{u}(s) - \mathbf{u}(t_{j+1})\|_{\frac{1}{2}+\beta}^2 ds, \\ \mathcal{N}_{j,1} := \int_{t_j}^{t_{j+1}} \|\mathbf{u}(s) - \mathbf{u}(t_{j+1})\|_\beta^2 (\|U^j\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2) ds, \\ \mathcal{N}_{j,2} := \int_{t_j}^{t_{j+1}} \|\mathbf{e}^{j+1}\|_\beta^2 (\|U^{j+1}\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2) ds, \\ \mathcal{N}_{j,3} := \int_{t_j}^{t_{j+1}} \|\mathbf{u}(s) - \mathbf{u}(t_j)\|_\beta^2 (|U^{j+1}|^2 + |\mathbf{u}(s)|^2) ds, \\ \mathcal{N}_{j,4} := \int_{t_j}^{t_{j+1}} \|\mathbf{e}^j\|_\beta^2 (|U^{j+1}|^2 + |\mathbf{u}(s)|^2) ds, \\ \mathcal{N}_{j,5} := \int_{t_j}^{t_{j+1}} \|(I - \pi_N)B(\mathbf{u}(s), \mathbf{u}(s))\|_{\beta-\frac{1}{2}}^2 ds, \\ \mathcal{W}_j := \int_{t_j}^{t_{j+1}} \langle A^{2\beta}\mathbf{e}^{j+1}, [G(\mathbf{u}(s)) - \pi_N G(U^j)] dW(s) \rangle.$$

Let  $m \in [1, M]$  an arbitrary integer. Summing (52) from  $j = 0$  to  $m - 1$ , multiplying by  $\mathbb{1}_{\Omega_k}$ , taking the mathematical expectation, and finally taking the maximum over  $m \in [1, M]$  imply

$$\begin{aligned} & \max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] + k \sum_{j=1}^M \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \\ & \leq \mathbb{E} \|\mathbf{e}^0\|_\beta^2 + 16C_0^2 \sum_{j=0}^{M-1} \sum_{i=1}^5 \mathbb{E} [\mathbb{1}_{\Omega_k} \mathcal{N}_{j,i}] + 2 \sum_{j=0}^{M-1} \mathbb{E} [\mathbb{1}_{\Omega_k} \mathcal{L}_j] + 2 \max_{1 \leq m \leq M} \sum_{j=0}^{m-1} \mathbb{E} [\mathbb{1}_{\Omega_k} \mathcal{W}_j]. \end{aligned}$$

Invoking the two items of Lemma 4.1 and the fact that  $\|\mathbf{u}(s)\|_\beta^2 + \max_{0 \leq j \leq M} \|\mathbf{U}^j\|_\beta^2 \leq f(k)$  on the set  $\Omega_k$  we infer that

$$\begin{aligned} & \max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] + k \sum_{j=1}^M \mathbb{E} \left[ \mathbb{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \\ (53) \quad & \leq \mathbb{E} \|\mathbf{e}^0\|_\beta^2 + 16C_0^2 k f(k) \sum_{j=0}^{M-1} \mathbb{E} \left( \mathbb{1}_{\Omega_k} [\|\mathbf{e}^{j+1}\|_\beta^2 + \|\mathbf{e}^j\|_\beta^2] \right) + 2C_8 f(k) M k [\Psi(k) + k^{1+\frac{1}{2}-\beta}] \\ & \quad + 64C_0^2 C_8 [f(k)]^2 M k [\Psi(k) + k] + 16C_0^2 \sum_{j=0}^{M-1} \mathcal{N}_{j,5} + 2 \max_{1 \leq m \leq M} \sum_{j=0}^{m-1} \mathbb{E} [\mathbb{1}_{\Omega_k} \mathcal{W}_j], \end{aligned}$$

where  $\psi(k) := k^{2-2\beta} + k^{2(\frac{1}{4}-\beta)}$ . Now, thanks to Assumption (B1)' we have

$$\begin{aligned} \mathbb{1}_{\Omega_k} \int_{t_j}^{t_{j+1}} \|(\mathbf{I} - \pi_N) \mathbf{B}(\mathbf{u}(s), \mathbf{u}(s))\|_{\beta-\frac{1}{2}}^2 ds &= \mathbb{1}_{\Omega_k} \int_{t_j}^{t_{j+1}} \sum_{n=N+1}^{\infty} \lambda_n^{2\beta-1} |\mathbf{B}_n(\mathbf{u}(s), \mathbf{u}(s))|^2 ds \\ &\leq \lambda_N^{2\beta-1} \int_{t_j}^{t_{j+1}} \mathbb{1}_{\Omega_k} \sum_{n=0}^{\infty} |\mathbf{B}_n(\mathbf{u}(s), \mathbf{u}(s))|^2 ds \\ &\leq \lambda_N^{2\beta-1} \int_{t_j}^{t_{j+1}} \mathbb{1}_{\Omega_k} |\mathbf{B}(\mathbf{u}(s), \mathbf{u}(s))|^2 ds \\ &\leq C \lambda_N^{2\beta-1} k \sup_{s \in [0, T]} \|\mathbf{u}(s)\|_{\frac{1}{4}}^4. \end{aligned}$$

Hence, owing to (15) we find a constant  $C > 0$  such that

$$\mathbb{E} \mathbb{1}_{\Omega_k} \int_{t_j}^{t_{j+1}} \|(\mathbf{I} - \pi_N) \mathbf{B}(\mathbf{u}(s), \mathbf{u}(s))\|_{\beta-\frac{1}{2}}^2 ds \leq C \lambda_N^{2\beta-1} k.$$

Notice also that

$$\begin{aligned} & \sum_{j=0}^{M-1} \|\mathbf{e}^{j+1}\|_\beta^2 (\|\mathbf{U}^{j+1}\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2) \\ &= \sum_{j=0}^{M-1} \|\mathbf{U}^{j+1} - \mathbf{U}^j + \mathbf{U}^j - \mathbf{u}(t_j) + \mathbf{u}(t_j) - \mathbf{u}(t_{j+1})\|_\beta^2 (\|\mathbf{U}^{j+1}\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2) \\ &\leq 3 \sum_{j=0}^{M-1} \left( \|\mathbf{U}^{j+1} - \mathbf{U}^j\|_\beta^2 + \|\mathbf{e}^j\|_\beta^2 + \|\mathbf{u}(t_j) - \mathbf{u}(t_{j+1})\|_\beta^2 \right) \left( \max_{0 \leq j \leq M} \|\mathbf{U}^{j+1}\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2 \right). \end{aligned}$$

Therefore,

$$\begin{aligned} & \mathbb{E} \left( \mathbb{1}_{\Omega_k} \sum_{j=0}^{M-1} \|\mathbf{e}^{j+1}\|_\beta^2 (\|\mathbf{U}^{j+1}\|_\beta^2 + \|\mathbf{u}(s)\|_\beta^2) \right) - C f(k) \mathbb{E} \sum_{m=0}^{M-1} \|\mathbf{e}^j\|_\beta^2 + f(k) C_7 [\psi(k) + k] \\ & \leq C \left( \mathbb{E} \left( \sum_{j=0}^{M-1} \|\mathbf{U}^{j+1} - \mathbf{U}^j\|_\beta^2 \right)^2 \right)^{\frac{1}{2}} \left( \mathbb{E} \max_{0 \leq j \leq M} \|\mathbf{U}^j\|_\beta^4 + \mathbb{E} \sup_{s \in [0, T]} \|\mathbf{u}(s)\|_\beta^4 \right)^{\frac{1}{2}}. \end{aligned}$$

As long as the initial data is concerned, we have

$$(54) \quad \mathbb{E} \|\mathbf{e}^0\|_\beta^2 = \|[\pi_N + (\mathbf{I} - \pi_N)]\mathbf{u}_0 - \pi_N\mathbf{u}_0\|_\beta^2$$

$$(55) \quad \leq \sum_{n=N+1}^{\infty} \lambda_n^{2(\beta-\frac{1}{4})} \lambda_N^{\frac{1}{2}} |\mathbf{u}_{0,n}|^2$$

$$(56) \quad \leq \lambda_N^{2(\beta-\frac{1}{4})} \|\mathbf{u}_0\|^2.$$

From all the above observations, (53), Assumption (B1)', (26)-(28) and (30) we infer that there exists a constant  $C_9 > 0$  such that

(57)

$$\begin{aligned} & \max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] + k \sum_{j=1}^M \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \\ & \leq C_9 f(k) [\Psi(k) + k^{1+\frac{1}{2}-\beta}] + C_9 f(k) [\Psi(k) + k] + C_9 \left( \lambda_N^{2\beta-1} + \lambda^{2(\beta-\frac{1}{4})} \right) + 2 \max_{1 \leq m \leq M} \sum_{j=0}^{m-1} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \mathcal{W}_j \right] \\ & \quad + C_9 k f(k) \sum_{m=0}^{M-1} \max_{1 \leq j \leq m} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_\beta^\beta \right] + 16C_0^2 k f(k) \max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right]. \end{aligned}$$

Now we deal with the term containing  $\mathcal{W}_j$ . After subtracting from  $\mathcal{W}_j$  the martingale  $M_0$  with mean zero defined by

$$M_0 = \int_{t_j}^{t_{j+1}} \langle A^\beta \mathbf{e}^{j+1}, A^\beta [G(\mathbf{u}(s)) - \pi_N G(\mathbf{U}^j)] dW(s) \rangle,$$

then taking the mathematical expectation, using the Young inequality and the Itô isometry give

$$\begin{aligned} \mathbb{E} \mathbf{1}_{\Omega_k} \mathcal{W}_j & \leq C \mathbb{E} \mathbf{1}_{\Omega_k} \left\| \int_{t_j}^{t_{j+1}} [G(\mathbf{u}(s)) - \pi_N G(\mathbf{U}^j)] dW(s) \right\|_\beta^2 + \frac{1}{4} \mathbb{E} \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \\ & \leq C \int_{t_j}^{t_{j+1}} \mathbb{E} \mathbf{1}_{\Omega_k} \|G(\mathbf{u}(s)) - \pi_N G(\mathbf{U}^j)\|_{\mathcal{L}(\mathcal{H}, V_\beta)}^2 ds + \frac{1}{4} \mathbb{E} \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \\ & \leq \sum_{i=1}^3 \mathbb{E} [\mathbf{1}_{\Omega_k} \mathcal{W}_{j,i}] + \frac{1}{4} \mathbb{E} \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2, \end{aligned}$$

where the first two symbols  $\mathcal{W}_{j,i}$ ,  $i \in \{1, 2\}$  satisfy the following equalities and inequalities

$$\begin{aligned} \mathbb{E} [\mathbf{1}_{\Omega_k} \mathcal{W}_{j,1}] & = C \int_{t_j}^{t_{j+1}} \mathbb{E} \mathbf{1}_{\Omega_k} \|\pi_N G(\mathbf{u}(s)) - \pi_N G(\mathbf{u}(t_j))\|_{\mathcal{L}(\mathcal{H}, V_\beta)}^2 ds \\ & \leq CC_3^2 \int_{t_j}^{t_{j+1}} \mathbb{E} \|\mathbf{u}(s) - \mathbf{u}(t_j)\|_\beta^2 ds \\ & \leq CC_3^2 C_7^2 k [k^{2-2\beta} + k^{2(\frac{1}{4}-\beta)} + k]; \end{aligned}$$

$$\begin{aligned} \mathbb{E} [\mathbf{1}_{\Omega_k} \mathcal{W}_{j,2}] & = C \int_{t_j}^{t_{j+1}} \mathbb{E} \mathbf{1}_{\Omega_k} \|\pi_N G(\mathbf{u}(t_j)) - \pi_N G(\mathbf{U}^j)\|_{\mathcal{L}(\mathcal{H}, V_\beta)}^2 ds \\ & \leq CC_3^2 k \mathbb{E} \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_\beta^2, \end{aligned}$$

where Lemma 4.1 was used to get the last line.

The third term  $\mathcal{W}_{j,3}$  satisfies

$$\begin{aligned} \mathbb{E}[\mathbf{1}_{\Omega_k} \mathcal{W}_{j,3}] &= \int_{t_j}^{t_{j+1}} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \|(\mathbf{I} - \pi_N)G(\mathbf{u}(s))\|_{\mathcal{L}(\mathcal{H}, V_\beta)}^2 \right) ds \\ &= \int_{t_j}^{t_{j+1}} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \sum_{n=N+1}^{\infty} \lambda_n^{2(\beta-\frac{1}{4})} \lambda_n^{\frac{1}{2}} \sup_{h \in \mathcal{H}, \|h\|_{\mathcal{H} \leq 1}} |G_n(\mathbf{u}(s))h|^2 \right) ds \\ &\leq \lambda_N^{2(\beta-\frac{1}{4})} \int_{t_j}^{t_{j+1}} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \sum_{n=1}^{\infty} \lambda_n^{\frac{1}{2}} \sup_{h \in \mathcal{H}, \|h\|_{\mathcal{H} \leq 1}} |G_n(\mathbf{u}(s))h|^2 \right) ds \\ &\leq \lambda_N^{2(\beta-\frac{1}{4})} k \mathbb{E} \left( \mathbf{1}_{\Omega_k} \sup_{s \in [0, T]} \|G(\mathbf{u}(s))\|_{\mathcal{L}(\mathcal{H}, V_{\frac{1}{4}})}^2 \right). \end{aligned}$$

Now, using Assumption (G) and the estimate (15) we infer that

$$\mathbb{E}[\mathbf{1}_{\Omega_k} \mathcal{W}_{j,3}] \leq CC_3^2 \lambda_N^{2(\beta-\frac{1}{4})} k,$$

for any  $j \in [0, M]$ . Thus, summing up we have obtained that

$$\begin{aligned} 2 \max_{1 \leq m \leq M} \sum_{j=0}^{m-1} \mathbb{E}[\mathbf{1}_{\Omega_k} \mathcal{W}_j] &\leq CC_3^2 C_7^2 T [\psi(k) + k] + CC_3^2 T \lambda_N^{2(\beta-\frac{1}{4})} \\ &+ CC_3^2 k \sum_{m=0}^{M-1} \max_{1 \leq j \leq m} \mathbb{E}[\mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_\beta^2] + \frac{1}{2} \sum_{m=0}^{M-1} \mathbb{E} \left( \mathbf{1}_{\Omega_k} \|\mathbf{e}^{m+1} - \mathbf{e}^m\|_\beta^2 \right). \end{aligned}$$

By plugging this last estimate into (53), we find a constant  $C_{10} > 0$  such that

$$\begin{aligned} &\max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] + 2k \sum_{j=1}^M \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \\ &\leq C_{10} f(k) [\Psi(k) + k + k^{1+\frac{1}{2}-\beta}] + C_{10} f(k) [\Psi(k) + k] + C_{10} \lambda_N^{2\beta-1} + C_{10} \lambda_N^{2(\beta-\frac{1}{4})} \\ &\quad + C_{10} k [f(k) + 1] \sum_{m=0}^{M-1} \max_{1 \leq j \leq m} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_\beta^2 \right]. \end{aligned}$$

Now, an application of the discrete Gronwall lemma yields

$$\begin{aligned} &\max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] + 2k \sum_{j=1}^M \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \\ &\leq \left( C_{10} f(k) [\Psi(k) + k + k^{1+\frac{1}{2}-\beta}] + C_{10} f(k) [\Psi(k) + k] + C_{10} \lambda_N^{2\beta-1} + C_{10} \lambda_N^{2(\beta-\frac{1}{4})} \right) e^{C_{10} T [f(k)+1]}. \end{aligned}$$

Since

$$\min\{k^{2-2\beta}, k^{1+\frac{1}{2}-\beta}, k^{2(\frac{1}{4}-\beta)}, k\} = k^{2(\frac{1}{4}-\beta)} \text{ and } \min\{\lambda_N^{2(\beta-\frac{1}{4})}, \lambda_N^{2\beta-1}\} = \lambda_N^{2(\beta-\frac{1}{4})},$$

for any  $\beta \in [0, \frac{1}{4})$ , and  $k^\varepsilon f(k) = k^\varepsilon \log k^{-\varepsilon} \leq \frac{1}{2}$ , then for any  $k > 0$  and  $\varepsilon \in (0, 2(\frac{1}{4} - \beta))$ , we derive that there exists a constant  $C > 0$  such that

$$\begin{aligned} (58) \quad &\max_{1 \leq m \leq M} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^m\|_\beta^2 \right] + \sum_{j=0}^{M-1} \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^{j+1} - \mathbf{e}^j\|_\beta^2 \right] \\ &+ 2k \sum_{j=1}^M \mathbb{E} \left[ \mathbf{1}_{\Omega_k} \|\mathbf{e}^j\|_{\frac{1}{2}+\beta}^2 \right] \leq C k^{-2\varepsilon} [k^{2(\frac{1}{4}-\beta)} + \lambda_N^{-2(\frac{1}{4}-\beta)}]. \end{aligned}$$

This estimate completes the proof of the Theorem 2.8.  $\square$

## 5. MOTIVATING EXAMPLES

In this section we give two examples of evolution equations to which we can apply our abstract result.

**5.1. Stochastic GOY and Sabra shell models.** The first examples we can take is the GOY and Sabra shell models. To describe this model let us denote by  $\mathbb{C}$  the field of complex numbers,  $\mathbb{C}^{\mathbb{N}}$  the set of all  $\mathbb{C}$ -valued sequences, and we set

$$H = \left\{ \mathbf{u} = (\mathbf{u}_n)_{n \in \mathbb{N}} \subset \mathbb{C}; \sum_{n=1}^{\infty} |\mathbf{u}_n|^2 < \infty \right\}.$$

Let  $k_0$  be a positive number and  $\lambda_n = k_0 2^n$  be a sequence of positive numbers. The space  $H$  is a separable Hilbert space when endowed with the scalar product defined by

$$\langle \mathbf{u}, \mathbf{v} \rangle = \sum_{k=1}^{\infty} \mathbf{u}_k \bar{\mathbf{v}}_k, \text{ for } \mathbf{u}, \mathbf{v} \in H,$$

where  $\bar{z}$  denotes the conjugate of any complex number  $z$ .

We define a linear map  $A$  with domain

$$D(A) = \left\{ \mathbf{u} \in H; \sum_{n=1}^{\infty} \lambda_n^4 |\mathbf{u}_n|^2 < \infty \right\},$$

by setting

$$A\mathbf{u} = (\lambda_n^2 \mathbf{u}_n)_{n \in \mathbb{N}}, \text{ for } \mathbf{u} \in D(A).$$

It is not hard to check that  $A$  is a self-adjoint and strictly positive operator. Moreover, the embedding  $D(A^\alpha) \subset D(A^{\alpha+\varepsilon})$  is compact for any  $\alpha \in \mathbb{R}$  and  $\varepsilon > 0$ . Thanks to this observation we can and will assume that there exists an orthonormal basis  $\{\psi_n; n \in \mathbb{N}\}$  of  $H$  such that

$$A\psi_n = \lambda_n \psi_n.$$

We can characterize the spaces  $D(A^\alpha)$ ,  $\alpha \in \mathbb{R}$  as follow

$$D(A^\alpha) = \left\{ \mathbf{u} = (\mathbf{u}_n)_{n \in \mathbb{N}} \subset \mathbb{C}; \sum_{n=1}^{\infty} \lambda_n^{4\alpha} |\mathbf{u}_n|^2 < \infty \right\}.$$

For any  $\alpha \in \mathbb{R}$  the space  $V_\alpha = D(A^\alpha)$  is a separable Hilbert space when equipped with the scalar product

$$(59) \quad ((\mathbf{u}, \mathbf{v}))_\alpha = \sum_{k=1}^{\infty} \lambda_k^{4\alpha} \mathbf{u}_k \bar{\mathbf{v}}_k, \text{ for } \mathbf{u}, \mathbf{v} \in V_\alpha.$$

The norm associated to this scalar product will be denoted by  $\|\mathbf{u}\|_\alpha$ ,  $\mathbf{u} \in V_\alpha$ . In what follows we set  $V = D(A^{\frac{1}{2}})$ .

Now, let  $\alpha_0 > \frac{1}{2}$  and  $\{\mathbf{w}_j; j \in \mathbb{N}\}$  be a sequence of mutually independent and identically distributed standard Brownian motions on filtered complete probability space  $\mathfrak{U} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$  satisfying the usual condition. We set

$$W(t) = \sum_{n=0}^{\infty} \lambda_n^{-\alpha_0} \mathbf{w}_n(t) \psi_n.$$

The process  $W$  defines a  $H$ -valued process with covariance  $A^{-2\alpha_0}$  which is of trace class. We also consider a Lipschitz map  $g : [0, \infty) \rightarrow \mathbb{R}$  such that  $|g(0)| < \infty$ . We define a map  $G : H \rightarrow \mathcal{L}(H, V_{\frac{1}{4}})$  defined by

$$G(u)h = g(\|u\|_0)h, \text{ for any } u \in H, h \in H.$$

This map satisfies Assumption **(G)**.

With the above notation, the stochastic evolution equation describing our randomly perturbed GOY and Sabra shell models is given by

$$(60) \quad \begin{cases} d\mathbf{u} = [A\mathbf{u} + B(\mathbf{u}, \mathbf{u})]dt + G(u)dW, \\ \mathbf{u}(0) = \mathbf{u}_0, \end{cases}$$

where  $B(\cdot, \cdot)$  is a bilinear map defined on  $V \times V$  taking values in the dual space  $V^*$ . More precisely, we assume that the nonlinear term

$$\begin{aligned} B : \mathbb{C}^{\mathbb{N}} \times \mathbb{C}^{\mathbb{N}} &\rightarrow \mathbb{C}^{\mathbb{N}}, \\ (\mathbf{u}, \mathbf{v}) &\mapsto B(\mathbf{u}, \mathbf{v}) = (b_1(\mathbf{u}, \mathbf{v}), \dots, b_n(\mathbf{u}, \mathbf{v}), \dots) \end{aligned}$$

for the GOY shell model (see [30]) is defined by

$$\begin{aligned} b_n(\mathbf{u}, \mathbf{v}) &:= (B(\mathbf{u}, \mathbf{v}))_n \\ &:= i\lambda_n \left( \frac{1}{4}\bar{v}_{n-1}\bar{u}_{n+1} - \frac{1}{2}(\bar{u}_{n+1}\bar{v}_{n+2} + \bar{v}_{n+1}\bar{u}_{n+2}) + \frac{1}{8}\bar{u}_{n-1}\bar{v}_{n-2} \right), \end{aligned}$$

and for the Sabra shell model, it is defined by

$$\begin{aligned} b_n(\mathbf{u}, \mathbf{v}) &:= (B(\mathbf{u}, \mathbf{v}))_n := \frac{i}{3}\lambda_{n+1} [\bar{v}_{n+1}u_{n+2} + 2\bar{u}_{n+1}v_{n+2}] \\ &\quad + \frac{i}{3}\lambda_n [\bar{u}_{n-1}v_{n+1} - \bar{v}_{n-1}u_{n+1}] \\ &\quad + \frac{i}{3}\lambda_{n-1} [2u_{n-1}v_{n-2} + u_{n-2}v_{n-1}], \end{aligned}$$

for any  $\mathbf{u} = (u_1, \dots, u_n, \dots) \in \mathbb{C}^{\mathbb{N}}$  and  $\mathbf{v} = (v_1, \dots, v_n, \dots) \in \mathbb{C}^{\mathbb{N}}$ .

**Lemma 5.1.** (a) For any non-negative numbers  $\alpha$  and  $\beta$  such that  $\alpha + \beta \in (0, \frac{1}{2}]$ , there exists a constant  $c_0 > 0$  such that

$$(61) \quad \|\mathbf{B}(\mathbf{u}, \mathbf{v})\|_{-\alpha} \leq c_0 \begin{cases} \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta} & \text{for any } \mathbf{u} \in V_{\frac{1}{2}-(\alpha+\beta)}, \mathbf{v} \in V_{\beta} \\ \|\mathbf{u}\|_{\beta} \|\mathbf{v}\|_{\frac{1}{2}-(\alpha+\beta)} & \text{for any } \mathbf{v} \in V_{\frac{1}{2}-(\alpha+\beta)}, \mathbf{u} \in V_{\beta}. \end{cases}$$

(b) For any  $\mathbf{u} \in H, \mathbf{v} \in V$

$$(62) \quad \langle \mathbf{B}(\mathbf{u}, \mathbf{v}), \mathbf{v} \rangle = 0.$$

*Proof.* The item (b) was proved in [19, Proposition 1], thus we omit its proof.

Item (a) can be viewed as a generalization of [19, Proposition 1]. We will just prove the latter item for the Sabra shell model since the proofs for the two models are very similar. Let  $\mathbf{u} \in V_{\frac{1}{2}-(\alpha+\beta)}$ ,  $\mathbf{v} \in V_{\beta}$ , and  $\mathbf{w} \in V_{\alpha}$  such that  $\|\mathbf{w}\|_{\alpha} \leq 1$ . We have

$$\begin{aligned} |\langle \mathbf{B}(\mathbf{u}, \mathbf{v}), \mathbf{w} \rangle| &= \left| \sum_{n=1}^{\infty} b_n(\mathbf{u}, \mathbf{v}) \bar{w}_n \right| \leq \sum_{n=1}^{\infty} |b_n(\mathbf{u}, \mathbf{v})| |\bar{w}_n| \\ &\leq \frac{1}{3} \sum_{n=1}^{\infty} \lambda_{n+1} (|\mathbf{u}_{n+1}| \cdot |\mathbf{v}_{n+2}| + |\mathbf{u}_{n+2}| \cdot |\mathbf{v}_{n+1}|) |\bar{w}_n| \\ &\quad + \frac{1}{3} \sum_{n=1}^{\infty} \lambda_n (|\mathbf{u}_{n-1}| \cdot |\mathbf{v}_{n+1}| + |\mathbf{u}_{n+1}| \cdot |\mathbf{v}_{n-1}|) |\bar{w}_n| \\ &\quad + \frac{1}{3} \sum_{n=1}^{\infty} \lambda_{n-1} (|\mathbf{u}_{n-1}| \cdot |\mathbf{v}_{n-2}| + |\mathbf{u}_{n-2}| \cdot |\mathbf{v}_{n-1}|) |\bar{w}_n| \\ &\leq I_1 + I_2 + I_3. \end{aligned}$$

For the term  $I_1$  we have

$$\begin{aligned} I_1 &\leq \frac{1}{3} \sum_{n=1}^{\infty} \lambda_{n+1} |\mathbf{u}_{n+1}| \cdot |\mathbf{v}_{n+2}| |\bar{w}_n| + \frac{1}{3} \sum_{n=1}^{\infty} \lambda_{n+1} |\mathbf{u}_{n+2}| \cdot |\mathbf{v}_{n+1}| |\bar{w}_n| \\ &\leq I_{1,1} + I_{1,2}. \end{aligned}$$



We will treat the term  $I_{1,1}$ . By Hölder's inequality we have

$$\begin{aligned} I_{1,1} &\leq \frac{1}{3} \sum_{n=1}^{\infty} k_0 2\lambda^{1-2\alpha} |\mathbf{u}_{n+1}| \cdot |\mathbf{v}_{n+2}| \lambda_n^{2\alpha} |\mathbf{w}_n| \\ &\leq \frac{2}{3} k_0 \left( \sum_{n=1}^{\infty} k_0 2\lambda_n^{2-4(\alpha+\beta)} |\mathbf{u}_{n+1}|^2 \lambda_n^{4\beta} |\mathbf{v}_{n+2}|^2 \right)^{\frac{1}{2}} \left( \sum_{n=1}^{\infty} \lambda_n^{4\alpha} |\mathbf{w}_n|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Since  $\|\mathbf{w}\|_{\alpha} \leq 1$  and  $\lambda_{n+p} = k_0^p 2^p \lambda_n$  we can find a constant  $C > 0$  depending only on  $\alpha, \beta$  and  $k_0$  such that

$$\begin{aligned} I_{1,1} &\leq C \left( \max_{k \in \mathbb{N}} \lambda_{n+1}^{2-4(\alpha+\beta)} |\mathbf{u}_{n+1}|^2 \right)^{\frac{1}{2}} \left( \sum_{n=1}^{\infty} \lambda_{n+2}^{4\beta} |\mathbf{v}_n|^2 \right)^{\frac{1}{2}} \\ &\leq C \left( \sum_{n=1}^{\frac{1}{2}} \lambda_{n+1}^{4[\frac{1}{2}-(\alpha+\beta)]} |\mathbf{u}_{n+1}|^2 \right)^{\frac{1}{2}} \left( \sum_{n=1}^{\infty} \lambda_{n+2}^{4\beta} |\mathbf{v}_n|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

from which we easily derive that

$$I_{1,1} \leq C \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta}.$$

One can use an analogous argument to show that

$$I_{1,2} \leq C \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta}.$$

Hence,

$$I_1 \leq C \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta}.$$

Using a similar argument we can also prove that for any non-negative numbers  $\alpha$  and  $\beta$  satisfying  $\alpha + \beta \in (0, \frac{1}{2}]$  there exists a constant  $C > 0$  such that

$$I_2 + I_3 \leq C \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta},$$

for any  $\mathbf{u} \in V_{\frac{1}{2}-(\alpha+\beta)}$  and  $\mathbf{v} \in V_{\beta}$ . Therefore, for any non-negative numbers  $\alpha$  and  $\beta$  satisfying  $\alpha + \beta \in (0, \frac{1}{2}]$  we can find a constant  $C > 0$  such that

$$\|\mathbf{B}(\mathbf{u}, \mathbf{v})\|_{-\alpha} \leq C \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta},$$

for any  $\mathbf{u} \in V_{\frac{1}{2}-(\alpha+\beta)}$  and  $\mathbf{v} \in V_{\beta}$ . Interchanging the role of  $\mathbf{u}$  and  $\mathbf{v}$  we obtain that for any two numbers  $\alpha$  and  $\beta$  as above there exists a positive constant  $C$  such that

$$\|\mathbf{B}(\mathbf{u}, \mathbf{v})\|_{-\alpha} \leq C \|\mathbf{v}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{u}\|_{\beta},$$

for any  $\mathbf{v} \in V_{\frac{1}{2}-(\alpha+\beta)}$  and  $\mathbf{u} \in V_{\beta}$ . Thus, we have just completed the proof of the lemma for the Sabra shell model. As we mentioned earlier, the case of the GOY model can be dealt with a similar argument.  $\square$

For more mathematical results related to shell models we refer to [3], [6], [7], and references therein.

**5.2. Stochastic nonlinear heat equation.** Let  $\mathcal{O}$  be a bounded domain of  $\mathbb{R}^d$ ,  $d = 1, 2$ . We assume that its boundary  $\partial\mathcal{O}$  is of class  $C^{\infty}$ . Throughout this section we will denote by  $H^{\theta}(\mathcal{O})$ ,  $\theta \in \mathbb{R}$ , the (fractional) Sobolev spaces as defined in [43] and  $H_0^1(\mathcal{O})$  be the space of functions  $\mathbf{u} \in H^1$  such that  $\mathbf{u}|_{\partial\mathcal{O}} = 0$ . In particular, we set  $H = L^2(\mathcal{O})$  and we denote its scalar product by  $(\cdot, \cdot)$ .

We define a continuous bilinear map  $\mathfrak{a} : H_0^1(\mathcal{O}) \times H_0^1(\mathcal{O}) \rightarrow \mathbb{R}$  by setting

$$\mathfrak{a}(\mathbf{u}, \mathbf{v}) = (\nabla \mathbf{u}, \nabla \mathbf{v}),$$

for any  $\mathbf{u}, \mathbf{v} \in H_0^1(\mathcal{O})$ . Thanks to the Riesz representation there exists a densely linear map  $A$  with domain  $D(A) \subset H$  such that

$$\langle A\mathbf{v}, \mathbf{u} \rangle = \mathfrak{a}(\mathbf{v}, \mathbf{u}),$$

for any  $\mathbf{u}, \mathbf{v} \in H_0^1(\mathcal{O})$ . It is well known that  $A$  is a self-adjoint and definite positive and its eigenfunctions  $\{\psi_n; n \in \mathbb{N}\} \subset C^{\infty}(\mathcal{O})$  form an orthonormal basis of  $H$ . The family of eigenvalues associated to

$\{\psi_n; n \in \mathbb{N}\}$  is denoted by  $\{\lambda_n; n \in \mathbb{N}\}$ . Observe that the asymptotic behaviour of the eigenvalues is given by  $\lambda_n \sim \lambda_1 n^{\frac{2}{d}}$ . For any  $\alpha \in \mathbb{R}$  we set  $V_\alpha = D(A^\alpha)$ , in particular we put  $V := D(A^{\frac{1}{2}})$ . We always understand that the norm in  $V_\alpha$  is denoted by  $\|\cdot\|_0$ .

Now, let  $\alpha_0 > \frac{d+1}{4}$  and  $\{\mathbf{w}_j; j \in \mathbb{N}\}$  be a sequence of mutually independent and identically distributed standard Brownian motions on filtered complete probability space  $\mathfrak{U} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$  satisfying the usual condition. We set

$$W(t) = \sum_{n=0}^{\infty} \lambda_n^{-\alpha_0} \mathbf{w}_n(t) \psi_n.$$

The process  $W$  defines a  $H$ -valued with covariance  $A^{-2\alpha_0}$  which is of trace class. We also consider a Lipschitz map  $g : [0, \infty) \rightarrow \mathbb{R}$  such that  $|g(0)| < \infty$ . We define a map  $G : H \rightarrow \mathcal{L}(H, V_{\frac{1}{4}})$  defined by

$$G(u)h = g(\|u\|_0)h, \text{ for any } u \in H, h \in H.$$

This map satisfies Assumption **(G)**.

The second example we can treat is the stochastic nonlinear heat equation

$$(63a) \quad d\mathbf{u} - [\Delta \mathbf{u} - |\mathbf{u}|\mathbf{u}]dt = g(\|\mathbf{u}\|_0)dW,$$

$$(63b) \quad \mathbf{u} = 0 \text{ on } \partial\mathcal{O},$$

$$(63c) \quad \mathbf{u}(0, x) = \mathbf{u}_0 \quad x \in \mathcal{O}.$$

This stochastic system can be rewritten as an abstract stochastic evolution equation

$$d\mathbf{u} + [A\mathbf{u} + B(\mathbf{u}, \mathbf{u})]dt = G(\mathbf{u})dW, \quad \mathbf{u}(0) = \mathbf{u}_0 \in H,$$

where  $A$  and  $G$  are defined as above and the  $D(A^{-\frac{1}{2}})$ -valued nonlinear map  $B$  is defined on  $H \times D(A^{\frac{1}{2}})$  or  $D(A^{\frac{1}{2}}) \times H$  by setting

$$B(\mathbf{u}, \mathbf{v}) = |\mathbf{u}|\mathbf{v},$$

for any  $(\mathbf{u}, \mathbf{v}) \in H \times D(A^{\frac{1}{2}})$  or  $(\mathbf{u}, \mathbf{v}) \in D(A^{\frac{1}{2}}) \times H$ . It is clear that

$$(64) \quad \langle A\mathbf{v} + B(\mathbf{u}, \mathbf{v}), \mathbf{v} \rangle \geq \|\mathbf{v}\|_{\frac{1}{2}}^2,$$

for any  $\mathbf{u}, \mathbf{v} \in V$ . Here we should note that thanks to the solution of Kato's square root problem in [2, Theorem 1], see also [34, Section 7], we have  $\|\mathbf{u}\|_{\frac{1}{2}} \simeq |\nabla \mathbf{u}|$  for any  $\mathbf{u} \in H_0^1(\mathcal{O})$ , i.e,  $V = H_0^1(\mathcal{O})$ .

Now we claim that for any numbers  $\alpha \in [0, \frac{1}{2})$  and  $\beta \in (0, \frac{1}{2})$  such that  $\alpha + \beta \in (0, \frac{1}{2})$ , there exists a constant  $c_0 > 0$  such that

$$(65) \quad \|B(\mathbf{u}, \mathbf{v})\|_{-\alpha} \leq c_0 \begin{cases} \|\mathbf{u}\|_{\frac{1}{2}-(\alpha+\beta)} \|\mathbf{v}\|_{\beta} & \text{for any } \mathbf{u} \in V_{\frac{1}{2}-(\alpha+\beta)}, \mathbf{v} \in V_{\beta} \\ \|\mathbf{u}\|_{\beta} \|\mathbf{v}\|_{\frac{1}{2}-(\alpha+\beta)} & \text{for any } \mathbf{v} \in V_{\frac{1}{2}-(\alpha+\beta)}, \mathbf{u} \in V_{\beta}, \end{cases}$$

and

$$(66) \quad \|B(\mathbf{u}, \mathbf{v})\|_{-\frac{1}{2}} \leq c_0 \|\mathbf{u}\|_{\frac{1}{4}} \|\mathbf{v}\|_{\frac{1}{4}} \quad \text{for any } \mathbf{v} \in V_{\frac{1}{4}}, \mathbf{u} \in V_{\frac{1}{4}}.$$

To prove these inequalities, let  $\beta > 0$  such that  $\alpha + \beta < \frac{1}{2}$ . Since

$$\left(\frac{1}{2} - \alpha\right) + \left(\frac{1}{2} - 1 + 2(\alpha + \beta)\right) + \left(\frac{1}{2} - \beta\right) = 1,$$

we have

$$(67) \quad |\langle |\mathbf{u}|\mathbf{v}, \mathbf{w} \rangle| \leq C_0 \|\mathbf{u}\|_{L^r} \|\mathbf{v}\|_{L^s} \|\mathbf{w}\|_{L^q},$$

where the constants  $q, r, s$  are defined through

$$\frac{1}{q} = \frac{1}{2} - \alpha, \quad \frac{1}{s} = \alpha + \beta, \quad \frac{1}{r} = \frac{1}{2} - \beta.$$

Recall that  $V_\alpha \subset H^{2\alpha} \subset L^q$  with  $\frac{1}{q} = \frac{1}{2} - \alpha$  if  $\alpha \in (0, \frac{1}{2})$  and  $q \in [2, \infty)$  arbitrary if  $\alpha = \frac{1}{2}$ . Then, we derive from (67) that the second inequality in (65) holds. By interchanging the role of  $r$  and  $s$  we

derive that the first inequality in (65) also holds. One can establish (66) with the same argument. The estimates (65) and (66) easily imply (3) and (7).

Now we need to check that  $B(\cdot, \cdot)$  satisfies (4). For this purpose we observe that there exists a constant  $C > 0$  such that

$$|B(\mathbf{u}, \mathbf{v})| \leq C \|\mathbf{u}\|_0 \|\mathbf{v}\|_{L^\infty},$$

which with the continuous embedding  $V_{\frac{1}{2}+\varepsilon} \subset L^\infty$  for any  $\varepsilon > 0$  implies (4).

## REFERENCES

- [1] A. Andersson and S. Larsson. Weak convergence for a spatial approximation of the nonlinear stochastic heat equation. *Math. Comp.*, 85(299):1335–1358, 2016.
- [2] P. Auscher and P. Tchamitchian. Square roots of elliptic second order divergence operators on strongly Lipschitz domains:  $L^2$  theory. *Journal d'Analyse Mathématique*, 90(1):1–12, 2003.
- [3] D. Barbato, M. Barsanti, H. Bessaih, and F. Flandoli. Some rigorous results on a stochastic GOY model. *Journal of Statistical Physics*, 125(3):677–716, 2006.
- [4] A. Bensoussan and R. Temam. Equations stochastiques du type Navier–Stokes. *Journal of Functional Analysis*, 13(2):195 – 222, 1973.
- [5] H. Bessaih, Z. Brzeźniak, and A. Millet. Splitting up method for the 2d stochastic Navier–Stokes equations. *Stochastic Partial Differential Equations: Analysis and Computations*, 2(4):433–470, 2014.
- [6] H. Bessaih and B. Ferrario. Invariant Gibbs measures of the energy for shell models of turbulence: the inviscid and viscous cases. *Nonlinearity*, 25(4):1075, 2012.
- [7] H. Bessaih, F. Flandoli, and E. Titi. Stochastic attractors for shell phenomenological models of turbulence. *Journal of Statistical Physics*, 140(4):688–717, 2010.
- [8] H. Bessaih and H. Schurz. Upper bounds on the rate of convergence of truncated stochastic infinite-dimensional differential systems with regular noise. *Journal of Computational and Applied Mathematics*, 208(2):354 – 361, 2007.
- [9] D. Blömker and A. Jentzen. Galerkin approximations for the stochastic Burgers equation. *SIAM J. Numer. Anal.* 51(1): 694–715, 2013.
- [10] H. Breckner. Galerkin approximation and the strong solution of the Navier-Stokes equation. *J. Appl. Math. Stochastic Anal.*, 13(3):239–259, 2000.
- [11] Z. Brzeźniak, E. Carelli, and A. Prohl. Finite-element-based discretizations of the incompressible Navier-Stokes equations with multiplicative random forcing. *IMA J. Numer. Anal.*, 33(3):771–824, 2013.
- [12] Z. Brzeźniak, E. Hausenblas, and P. Razafimandimby. Stochastic nonparabolic dissipative systems modeling the flow of liquid crystals : Strong solution (mathematical analysis of incompressible flow). *RIMS Kokyuroku*, 1875:41–72, jan 2014.
- [13] Z. Brzeźniak and E. Motyl. Existence of a martingale solution of the stochastic Navier–Stokes equations in unbounded 2D and 3D domains. *Journal of Differential Equations*, 254(4):1627 – 1685, 2013.
- [14] T. Caraballo, J. Real, and T. Taniguchi. On the existence and uniqueness of solutions to stochastic three-dimensional lagrangian averaged Navier–Stokes equations. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 462(2066):459–479, 2006.
- [15] E. Carelli, E. Hausenblas and A. Prohl. Time-splitting methods to solve the stochastic incompressible Stokes equation. *SIAM, J. Numer. Anal.* 50(6): 2917–2939, 2012.
- [16] E. Carelli, A. Müller, and A. Prohl. Domain decomposition strategies for the stochastic heat equation. *Int. J. Comput. Math.*, 89(18):2517–2542, 2012.
- [17] E. Carelli and A. Prohl. Rates of convergence for discretizations of the stochastic incompressible Navier–Stokes equations. *SIAM J. Numer. Anal.*, 50(5):2467–2496, 2012.
- [18] P-L. Chow. *Stochastic Partial Differential Equations*, Chapman & Hall/CRC. Boca Raton, Florida, 2007.
- [19] P. Constantin, B. Levant, and E. S. Titi. Analytic study of shell models of turbulence. *Physica D: Nonlinear Phenomena*, 219(2):120 – 141, 2006.
- [20] J. Cui, J. Hong and Z. Liu. Strong convergence rate of finite difference approximations for stochastic cubic Schrödinger equations. *em Journal of Differential Equations*. 263(7): 3687–3713, 2017.
- [21] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2014.
- [22] R. Dautray and J. Lions. *Mathematical Analysis and Numerical Methods for Science and Technology: Volume 5 Evolution Problems I*. 1999.
- [23] A. de Bouard and A. Debussche. A semi-discrete scheme for the stochastic nonlinear Schrödinger equation. *Numerische Mathematik*, 96(4):733–770, 2004.
- [24] A. Debussche and J. Printems. Weak order for the discretization of the stochastic heat equation. *Math. Comp.*, 78(266):845–863, 2009.
- [25] A. Debussche and J. Printems. Convergence of a semi-discrete scheme for the stochastic Korteweg-de Vries equation, *Discrete and Continuous Dynamical Systems, series B*, 6(4): 761–781, 2006.

- [26] A. Debussche and J. Printems. Numerical simulation of the stochastic Korteweg-de Vries equation, *Physica D: Nonlinear Phenomena*, 134: 200–226, 1999.
- [27] P. D. Ditlevsen. *Turbulence and Shell Models*. Cambridge University Press, Cambridge, 2011.
- [28] P. Dörsek. Semigroup splitting and cubature approximations for the stochastic Navier-Stokes equations. *SIAM J. Numer. Anal.* 50(2): 729–746, 2012.
- [29] E. Gkioulekas. Review of ‘turbulence and shell models’ by Peter D. Ditlevsen. *SIAM Rev.*, 54(2):396–399, 2012.
- [30] E. B. Gledzer. System of hydrodynamic type admitting two quadratic integrals of motion. *Soviet Physics Doklady*, 18:216, Oct. 1973. Provided by the SAO/NASA Astrophysics Data System.
- [31] I. Gyöngy and A. Millet. Rate of convergence of space time approximations for stochastic evolution equations. *Potential Analysis*, 30(1):29–64, 2009.
- [32] M. Hairer. An introduction to stochastic PDEs. *arXiv preprint arXiv:0907.4178*, 2009.
- [33] A. Jentzen and P. E. Kloeden. The numerical approximation of stochastic partial differential equations. *Milan J. Math.* 77: 205–244, 2009.
- [34] D. Jerison and C. E. Kenig, The inhomogeneous Dirichlet problem in Lipschitz domains, *J. Funct. Anal.* 130(1): 161–219, 1995.
- [35] A. N. Kolmogorov. The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Proceedings: Mathematical and Physical Sciences*, 434(1890):9–13, 1991.
- [36] N. V. Krylov. A relatively short proof of Itô’s formula for SPDEs and its applications. *Stoch. Partial Differ. Equ. Anal. Comput.*, 1(1):152–174, 2013.
- [37] N. V. Krylov and B. L. Rozovskii. Stochastic evolution equations. *Journal of Soviet Mathematics*, 16(4): 1233–1277, 1981.
- [38] F. Lindner and R. L. Schilling. Weak order for the discretization of the stochastic heat equation driven by impulsive noise. *Potential Anal.*, 38(2):345–379, 2013.
- [39] F. Plunian, R. Stepanov, and P. Frick. Shell models of magnetohydrodynamic turbulence. *Physics Reports*, 523(1):1 – 60, 2013.
- [40] C. Prévôt and M. Röckner. *A concise course on stochastic partial differential equations, Lecture Notes in Mathematics*. Berlin-Heidelberg-New York: Springer, 2007.
- [41] J. Printems. On the discretization in time of parabolic stochastic partial differential equations. *M2AN. Math. Model. Numer. Anal.* 35(6): 1055–1078, 2001.
- [42] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion.*, Springer, Berlin, 1994.
- [43] H. Triebel. *Interpolation Theory - Function Spaces - Differential Operators*. Wiley, 1999.

(Hakima Bessaih) UNIVERSITY OF WYOMING, DEPARTMENT OF MATHEMATICS, LARAMIE, WY 82071-3036, UNITED-STATES

(Erika Hausenblas) MONTANUNIVERSITÄT LEOBEN, LEHRSTUHL FÜR ANGEWANDTE MATHEMATIK, PETER-TUNNER-STRASSE 25-27, 8700 LEOBEN, AUSTRIA

(Tsiry Randrianasolo) BIELEFELD UNIVERSITY, DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITÄTSSTRASSE 25, 33615 BIELEFELD, GERMANY

*E-mail address*, Corresponding author: [trandria@math.uni-bielefeld.de](mailto:trandria@math.uni-bielefeld.de)

(Paul Razafimandimby) DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF PRETORIA, LYNWOOD ROAD, HATFIELD, PRETORIA 0083, SOUTH AFRICA